


No. 1388

1388
1877

Library Of
JAMES P. MARSH.



Longacre

BOSTON MEDICAL LIBRARY
in the Francis A. Countway
Library of Medicine ~ *Boston*

Digitized by the Internet Archive
in 2011 with funding from
Open Knowledge Commons and Harvard Medical School

<http://www.archive.org/details/contributionstor00buck>

CONTRIBUTIONS

TO

REPARATIVE SURGERY:

SHOWING

ITS APPLICATION TO THE TREATMENT OF DEFORMITIES, PRODUCED BY DESTRUCTIVE DISEASE OR INJURY; CONGENITAL DEFECTS FROM ARREST OR EXCESS OF DEVELOPMENT; AND CICATRICIAL CONTRACTIONS FROM BURNS.

BY

GURDON BUCK, M. D.

ILLUSTRATED BY NUMEROUS ENGRAVINGS.

NEW YORK:
D. APPLETON AND COMPANY,
549 AND 551 BROADWAY.
1876.

ENTERED, according to Act of Congress, in the year 1876,
BY D. APPLETON & CO.,
In the Office of the Librarian of Congress, at Washington.

P R E F A C E .

THE volume now offered to the medical profession contains the author's own experience in the department of practical surgery of which it treats. While it makes no claim to be a systematic and thorough treatise on the subject, it will, nevertheless, be found to contain, in a classified order, much that is treated in existing works on plastic surgery. It is believed also, that, in addition to this, something new and useful has been contributed to the resources of surgical art. There is no department of surgery where the ingenuity and skill of the surgeon are more severely taxed than when required to repair the damage sustained by the loss of parts, or to remove the disfigurement produced by destructive disease or violence, or to remedy the deformities of congenital malformation. The results obtained in such cases within the last half century are among the

most satisfactory achievements of modern surgery. The term "Reparative Surgery," chosen as the title of this volume, though it may, in a comprehensive sense, be applied to the treatment of a great variety of lesions to which the body is liable, is, however, restricted in this work exclusively to what has fallen under the author's own observation, and been subjected to the test of experience in his own practice. It largely embraces the treatment of lesions of the face, a region in which plastic surgery finds its most frequent and important application. Another and no less important class of lesions will also be found to have occupied a large share of the author's attention, viz., cicatricial contractions following burns. While these cases have a very strong claim upon our commiseration, and should stimulate us, as surgeons, to the greatest efforts for their relief, they have too often in the past been dismissed as hopelessly incurable. The satisfactory results obtained in the cases reported in this volume will encourage other surgeons, we trust, to resort with greater hopefulness in the future to operative interference. Accuracy of description and clearness of statement have been aimed at in the following pages; and if, in his endeavor to attain this impor-

tant end, the author has incurred the reproach of tediousness, the difficulty of the task must be his apology.

The indispensable aid of pictorial illustration has been employed to the fullest possible extent; and whatever success has been achieved in this department, the credit of it is due to the admirable skill of Ferdinand Froning, of Vienna, who executed all the principal figures from photographs. Special pains have also been taken to ascertain the permanent results of all the cases reported, after the lapse of long intervals of time, and fortunately this has been practicable in almost all instances.

Two albums, containing each a collection of the photographs from which the portrait figures illustrating this volume were engraved, have been deposited, the one in the Pathological Museum of the New York Hospital, the other in the United States Army Medical Museum, at Washington, D. C., where they are accessible to any person desirous of consulting them. Each photograph is designated by a number corresponding to the figure in the volume of which it is the original.

Of the cases reported in this volume, the following have previously appeared in print:

- CASE I. In "Transactions of the Medical Society of the State of New York," vol. 1864.
- II. "Transactions of the American Medical Association," vol. 1870.
- IV. *American Practitioner*, August, 1873.
- VI. *American Journal of Medical Science*, vol. lviii., new series, 1869, p. 352.
- VIII. "Transactions of the Medical Society of the State of New York," vol. 1866.
- IX. *New York Medical Record*, February 1, 1871.
- X. *Boston Medical and Surgical Journal*, January, 1874.
- XI. XII., XIII., XIV. *New York Medical Record*, January 15, 1872.
- XVII. *New York Medical Record*, 1866.
- XXI. *New York Medical Record*, 1870, p. 483.
- XXII. "Transactions of the New York State Medical Society," 1872.
- XXV. *New York Medical Record*, 1869.
- XXVII. *American Journal of Medical Science*, 1872, p. 52.
- XXVIII. *New York Medical Record*, 1874, p. 473.

C O N T E N T S.

CHAP.	PAGE
I. TRANSPLANTATION OF SKIN	7
II. METHODS OF TRANSFER	10
III. TREATMENT OF RAW SURFACES LEFT TO HEAL BY GRANULATION	13
IV. SUTURES AND THEIR MANAGEMENT	14
V. METHODS OF OPERATION	20

C A S E S.

FIRST CLASS.—LOSS OF PARTS INVOLVING THE FACE, AND RESULTING FROM DESTRUCTIVE DISEASE OR INJURY.

CASE I. Reconstruction of the Mouth and Repair of the Nose after the Loss of the Right Half of the Upper Lip, the Adjacent Portion of the Cheek and Ala Nasi, together with the Entire Right Superior Maxillary Bone	31
II. Reconstruction of the Mouth and Repair of the Nose after the Loss of the Right Half of the Upper Lip, and Adjacent Portion of the Cheek, and Right Ala Nasi	52
III. Reconstruction of the Nose and Mouth after the Loss of the Nose and Entire Upper Lip	62
IV. Reconstruction of the Mouth after the Loss of the Entire Under Lip, and a Portion of the Inferior Maxillary Bone	75
V. Reconstruction of the Mouth after Removal of the Under Lip for Disease	87
VI. Reconstruction of the Mouth after the Loss of the Right Half of Both Lips; also, Esmarch's Operation for Ankylosis of the Jaw	93
VII. Mutilation and Distortion of the Mouth from a Shell-Wound	101
VIII. Loss of a Large Portion of the Lower Jaw-bone, with Extensive Mutilation of the Face and Distortion of the Mouth, produced by a Shell-Wound	108
IX. Closure of an Opening into the Superior Meatus of the Right Nasal Fossa	114
X. A Rhinoplastic Operation for the Restoration of the Apex Nasi after it had been bitten off	118

SECOND CLASS.—CONGENITAL DEFECTS FROM ARREST OR EXCESS OF DEVELOPMENT.

—HARELIP.

	PAGE
CASE XI. Single Harelip	131
XII. Double Harelip	132
XIII. Single Harelip, with Cleft of the Dental Arch and Bony and Soft Palates	134
XIV. Single Harelip on the Right Side	137
XV. Single Harelip	138
XVI. Single Harelip	139
XVII. Double Harelip, complicated with Cleft of the Bony and Soft Palates, together with the Presence of an Intermaxillary Bone	140
XVIII., XIX., XX. Examples of Cases of Harelip, operated on a Second Time for the Purpose of remedying the Imperfect Results of Previous Operations performed in Infancy	144-146
XXI. Congenital Hypertrophy of the Tongue	151
XXII. Congenital Hypertrophy of the Under Lip	159
XXIII. Abnormal Growth of Hair upon the Forehead	165
XXIV. Erectile Tumor of Large Size	168
XXV. A Pendulous Tumor (Molluscum Fibrosum) arising from the Right Half of the Forehead and Temple	172

THIRD CLASS.—CICATRICIAL CONTRACTIONS FOLLOWING BURNS.

XXVI. Disfigurement of the Face from Cicatricial Contractions; Extirpa- tion of one Eyeball; Closure of the Orbit by a Plastic Opera- tion	185
XXVII. Cicatricial Contractions involving the Chin and Front of the Neck	197
XXVIII. Cicatricial Contractions involving the Face and Hand	206
XXIX. Cicatricial Contractions involving the Right Axilla and Arm	228

REPARATIVE SURGERY.

To avoid repetition in the succeeding narrative of cases, some general observations on the subjects designated by the titles of the following chapters will now be presented.

CHAPTER I.

TRANSPLANTATION OF SKIN.

Choice of Material.—In the choice of skin, which is the material employed to supply an existing deficiency, care should be taken that it be in a normal and healthy condition. If, in consequence of a previous burn, the surface has become glossy, pale, and of cicatricial formation, a patch taken from it for transplantation, by dissecting it up from its underlying connections, would inevitably slough. Such a patch could not maintain its vitality if nourished by the circulation it would receive through its pedicle alone; the vascular support derived from its connec-

tion with underlying parts could not be dispensed with. The author's experience on this point conclusively establishes the fact that cicatricial integument cannot be relied upon for transplantation, and should not be used for that purpose. Another condition to be observed is, that a patch of skin intended for transfer should have its long axis correspond to the direction in which the arterial vessels are distributed, and the free extremity of the patch should point toward their destination. Upon the forehead, for example, these important conditions are best secured by raising the patch of skin required from above the inner half of the eyebrow, where it would retain its connection and derive its vascular support from a branch of the ophthalmic artery, which emerges from the orbit through the supra-orbital notch.

Precision in Adaptation.—To secure precision in adapting a patch of skin to a new locality to which it is to be transferred, the following method will be found satisfactory. After having prepared the space which it is intended to fill up, by paring its edges, and dissecting them up sufficiently from their underlying connections to permit them to be everted, and also, if necessary, by notching the edges at intervals to facilitate their eversion, an exact pattern of the space should be cut from oiled silk, and the pattern applied to the surface which is to supply the new material. Small pins may then be temporarily in

served erect in the skin at intervals around the pattern, and at a distance of one line from its margin, as an allowance for shrinkage. A larger allowance, however, must be made for the length of the patch of skin beyond that of the pattern itself, so as to permit the patch to be brought around edgewise to its new location, without causing any strain at its pedicle, which might obstruct the circulation, and thereby endanger the vitality of the patch. The outline of the pattern having been indicated by the pins, the pattern itself may be dispensed with, and the pins alone left to serve as a guide. After the incision defining the patch of skin has been made, the patch itself should be dissected up from its underlying connections. Special care should be taken on the forehead not to wound or detach the pericranium.

CHAPTER II.

METHODS OF TRANSFER.

THE transfer of a portion of skin to a neighboring locality may be made by different methods, the choice of them depending upon the condition of the parts involved. In the cases hereinafter reported the methods employed were the following:

First Method. By Approximation.—If on both sides of a space requiring to be filled up the adjoining skin is supple and movable, the opposite edges of the space may be pared, and the adjacent skin dissected up from its underlying connections to a sufficient distance to permit the edges to meet and be secured in contact by sutures. If there be any strain on the sutures after the new adjustment, it should be relieved by an incision through the skin on either side of the wound, parallel with, and at a suitable distance from it.

Second Method. By Sliding.—If, upon one side only of a space requiring to be covered, the skin is in a condition to be made available for transfer, a patch of the required size, adjacent to the space, may then be dissected up, and, being left connected at

one end, may be glided edgewise toward the space, and adjusted to it by sutures; the edges of the space must of course have been prepared beforehand for the purpose. The surface still left bare by the transferred patch may sometimes be covered by dissecting up the neighboring skin, and stretching it across the bare surface till it meets the edge of the patch, where it may be adjusted by sutures.

Third Method. By Transfer to a Distance.— Sometimes the material for supplying a deficiency has to be taken from a locality at a distance from the spot where the deficiency exists. This may be done in two ways: 1. A patch of skin, after having been dissected up from its underlying connections, but still remaining attached at one extremity, may be transferred edgewise, and made to describe in its circuit a quadrant of a circle, or even an entire semi-circle, in reaching its new destination. In order that the raw under-surface of the patch, when thus transferred, may lie in contact throughout its entire length with subjacent raw surface, the intervening skin must be displaced. This displacement, however, should be effected in such a way that the displaced skin, retaining a connecting pedicle for its support, may be made to change places with the transplanted patch, and thus contribute, as far as it can, a covering for the surface that has been left bare. This method has been advantageously employed in several of the

cases reported in this volume. In all of them the forehead was the seat of operation, and in no instance has a patch of skin, taken from the forehead in the manner already described, failed to do well from deficient vitality, or sustained any loss by sloughing.

2. The transfer of a patch of skin to a distant spot may be effected by a jumping process, as follows: An elongated patch of skin having been dissected up, but left attached at one extremity, may be made to jump over an intervening sound surface, and have its free extremity adjusted to a distant spot where a deficiency exists. After becoming ingrafted in its new locality, the patch is to be severed, and its pedicle replaced upon the bare surface from which it was originally taken. But one example of this method occurs in the following pages (Case III., p. 62).

CHAPTER III.

TREATMENT OF RAW SURFACES LEFT TO HEAL BY GRANULATION.

AFTER the transfer of a patch of skin to a new locality, more or less of the surface from which it was taken must be left to heal by granulation. The following method of treating such a surface has proved uniformly satisfactory, and is therefore confidently recommended. After all hæmorrhage has ceased, the raw surface should be coated first with a uniform layer of scraped lint, and then with an additional layer of lint saturated with collodion. This dressing soon stiffens and forms an artificial scab, which will remain adherent for from six to ten days, when it becomes detached by suppuration. In the meantime the wound requires no other application; it remains in a quiescent state, with scarcely any surrounding redness or inflammatory tumefaction, and only a slight discharge of matter escapes from one or more points at the margin of the crust. On its separation a growth of healthy granulations will be found at the margin of the sore, and sometimes covering its entire surface, even up to the level of the surrounding skin. The application just described will hereafter be known and referred to as the collodion crust.

CHAPTER IV.

SUTURES AND THEIR MANAGEMENT.

IN plastic operations success depends so much on the proper management of sutures, that a particular description of the different methods of employing them will not be out of place. Three kinds of sutures have been employed by the author.

1. The Interrupted Thread Suture.—For the insertion of sutures of this sort, trocar-pointed needles, such as are used by glove-makers, are preferable to any other. They may be had of all sizes at the thread-and-needle stores. To guard against inversion of the edges of a wound, which is a frequent cause of the failure of primary union, the needle should be inserted in such a manner that it will pass obliquely through the thickness of the skin, and in doing so the deepest part of its track will be farther from the confronted edges of the wound than it is at the points of entrance and exit upon the surface. Thus inserted, the thread on being tightened tends to evert the edges of the wound, and bring their confronted cut surfaces more perfectly into contact. The confronting of the edges is further facilitated if,

after inserting the first suture, instead of tying the knot, the ends of the thread are made use of to draw out the edges of the wound, while a second neighboring suture is being inserted and tied; the same may be done with each additional suture in succession. The sutures should also be inserted as closely together as may be necessary to secure exact coaptation of the edges. Their multiplicity is not objectionable, inasmuch as at the expiration of twenty-four hours, when agglutination of the edges will have taken place, their number may be diminished by removing the alternate ones. Metallic sutures, in the judgment of the author, possess no advantage over thread, and are as liable to cause ulceration as thread. Both will remain harmless as long as they are needed, provided the edges of the wound they are designed to hold in contact have previously been liberated, so as to be relieved of all strain upon them.

2. The Pin, or Figure-of-eight, Suture.—The insertion of the pin is performed, with facility and precision, by the aid of an instrument devised by the author, and first described in the *New York Medical Record*, of July 1, 1869, under the name of "Suture Pin Conductor." It consists of a needle two and a half inches long, of the thickness of an ordinary knitting-needle, slightly curved toward the point, and fixed in a handle. From its point toward the

handle, its thickness grows smaller for a distance of half an inch, which facilitates its passage through the skin and beyond. Its extremity is beveled off to a sharp point on the concave side of the needle, and is perforated lengthwise for a short distance on its beveled face, like the point of an hypodermic syringe. (*See Fig. 1, a.*) It is used in the following manner: The edges of the wound which are to be approximated, having been traversed by the conductor guided by one hand, a pin, held between the thumb and fingers of the other hand, is engaged by




FIG. 1.

its point in the perforated hole at the point of the conductor, and held steadily in place while the conductor is withdrawn. (*Fig. 1, b.*) The pin follows it with perfect certainty to its place, and is then wound with loosely-twisted cotton-yarn passed in figure-of-eight turns around both ends of the pin.

The advantage of this instrument is, that with it the pin can be inserted with great precision, and fresh pins, when necessary, can be inserted near to and to supply the place of old ones requiring to be removed, without disturbing the newly-formed adhesions. To guard against suppuration in the track of the pin, and on the surface of the skin underlying the yarn, the yarn itself should be removed at the end of forty-eight hours, and sometimes of twenty-four hours, so as to allow the constricted surface to recover itself; after which fresh yarn should be applied. This change should be repeated daily afterward, till the removal of the pin on the fourth day, beyond which time it should not be continued in place. If the support afforded by the pin cannot yet be dispensed with on the fourth day, it is better to insert a fresh pin near by than to leave the old one in place longer than four days. A patch of three or four thicknesses of adhesive plaster stuck together, and cut of a proper shape, may be laid upon the surface of the skin, between the points of entrance and exit of the pin, for the yarn to rest upon, and to serve as an additional protection against ulceration.

3. The Beaded Wire Clamp Suture.—This suture is intended as an auxiliary for the support of other sutures, and may be employed where it is important to relieve them of strain, and thereby increase the

chances of obtaining primary union. The following is the method of inserting it. An ordinary darning-needle, two inches or more in length, may serve for the purpose, and should be threaded with a flexible silver wire, previously charged at its knotted extremity, first, with a small perforated disk of leather (such as may be found at any saddler's shop), and then with a smooth, round glass bead. The needle thus armed is then made to traverse the edges of the wound at opposite points, and at a distance of one inch or more from the edges. A second bead




FIG. 2.

should then be strung upon the other end of the wire. While the opposite edges of the wound are crowded toward each other and held in contact, the free end of the wire should be drawn upon, and the second bead slid down to its place against the skin, and secured there by winding the wire three or four times around the end of a friction match, or other suitable piece of wood. The wire itself should be left two inches long, so as to permit the clamp to be tightened or loosened at pleasure.

The further adjustment of the edges of the wound to each other may then be completed by any required number of interrupted thread sutures. The special advantage of the clamp suture is, that it relieves the other sutures of all strain, and admits of being tightened or slackened, if necessary. It may remain *in situ*, undisturbed, from six to ten days, and any superficial ulceration produced by the pressure of the beads against the skin heals readily after their removal, without leaving any scar. Another advantage is, that, if primary union wholly or partially fails to take place, the edges of the wound are still maintained in contact, while union by the slower process of granulation is going on.

CHAPTER V.

METHODS OF OPERATION.

THE observations under this head relate only to the reconstruction of the mouth after the entire loss of either the upper or lower lip, or of a considerable portion of either. The two methods about to be described have been successfully employed in several of the following cases, and, in the results obtained, it is believed these methods possess advantages over other methods in use among surgeons, and they also differ from them in some important respects. To avoid repetition, they will now be described in detail.

First Method.—In order to execute this method, a preliminary operation must be performed to bring the parts into a proper condition for it. For example, the under lip being involved, the removal of the diseased portion may be executed in two ways, the choice of which should be determined by the condition of the parts.

First Preliminary Operation is that for the removal of the diseased part, by including it in a let-

ter-V-shaped patch, as follows: An incision, commencing at a point within half an inch of the angle of the mouth on both sides and dividing the lip border should be carried downward on either side of the morbid growth in converging lines, till both incisions meet under the chin in the median plane. The included V-shaped patch should then be dissected up from the underlying periosteum and removed. The lining mucous membrane of the mouth should then be divided on both sides of the wound, along the line of its reflection from the jaw to the inside of the lip, and the division continued outward as far as is necessary to permit the edges of the V-shaped wound to meet at the symphysis, and there be secured together by sutures.

Second Preliminary Operation is that for the removal of the diseased part by including it in a quadrilateral patch. Two incisions, dividing the lip border at a distance of half an inch from the angles of the mouth, should extend vertically downward on either side of the diseased part, till they both connect with a third transverse incision crossing the lower part of the chin. After the removal of the included quadrilateral patch, the transverse incision should be continued outward, on both sides, to within a finger's breadth of the angles of the jaw, and thence upward a distance of two inches on the sides of the cheeks in lines curving forward. The two cheek-flaps thus

formed, both of which are lined with mucous membrane, are then to be dissected up from the jaw, and brought forward edgewise, so as to meet in front at the symphysis, where they are to be secured together by sutures. The surfaces left bare on both cheeks by this transfer may be covered again by dissecting up the skin bounding them posteriorly from its underlying connections, and advancing it so as to meet the edge of the transferred cheek-flap and be adjusted to it by sutures. By both these preliminary operations the mouth is made to assume a shape in which the upper lip is folded upon itself, and overhangs the retracted and shortened under lip. Case V., p. 87, furnishes an example of the application of the second method just described. Fig. 3 illustrates the first method as completed on the cadaver. The letter-V-shaped piece, excised from the under lip, remains attached and suspended below the chin, and the edges of the wound are secured together by sutures. After the parts thus adjusted have completely healed and regained their suppleness, the special operative method by which the mouth is to be restored to its natural shape may be undertaken. It is as follows. In order to insure precision in making the requisite incisions, their course should first be designated by pins temporarily inserted erect in the skin at certain points, as shown by Fig. 3. For example, letters *a a* represent two pins inserted

at one finger's breadth below the under-lip border, one on either side of the chin, a little to the outside of the angle of the mouth, and both equidistant from the median line; *b b* are also two pins inserted, one on either side, into the upper lip at the margin




FIG. 3.

of the vermilion border, both equidistant from the median line, and at such a distance apart as to include between them sufficient length of lip border with which to form a new upper lip. The steps of the operation are then the following: With the forefinger of the left hand placed on the inside of the mouth, the left cheek is to be kept moderately on the stretch, while with a sharp-pointed "Beer's" cor-

nea knife it is transfixed at the point *a*, marked by the lower pin on the left side of the chin. An incision is then to be carried through the entire thickness of the cheek, upward and a little outward, a distance of one inch and a half, to a point, *c*, near the middle of the cheek. The left half of the upper lip should next be transfixed at the point *b*, marked by a pin on the vermilion border, and the incision carried through the lip and cheek, outward and a little upward, to join the first incision at its terminus *c*, in the middle of the cheek. A triangular patch, *b, c, a*, will thus be formed, which will include the entire thickness of the cheek, and whose apex will be free and disconnected, while its base remains attached toward the mouth. The next step is to transfer this patch from the cheek to the side of the chin. For this purpose an incision should be made on the side of the chin from *a*, the starting-point of the first incision, vertically downward to the edge of the jaw, and to the depth of the periosteum. The edges of this incision retracting wide apart, afford a V-shaped space for the lodgment of the triangular patch, which is now to be brought around edgewise, and adjusted by sutures in its new locality on the side of the chin. By this transfer the portion of upper-lip border that formed a part of the base of the patch is brought into a transverse line, continuous with the under lip, and constitutes an extension of it. The space upon

the cheek from which the triangular patch was taken is closed by bringing its edges together, and securing them in contact by sutures. By this adjustment a new and naturally-shaped angle is formed for the mouth at the point *b*, on the vermillion border, where the lip was transfixed in commencing the second in-



FIG. 4.

cision of the cheek. The incisions described above should be made with the utmost precision, and special care should be taken that the lining mucous membrane is divided exactly to the same extent as the skin itself. The same procedure may be applied to the other side of the mouth and executed at the same operation, as was done in Case V., p. 87. The

mouth, when thus reconstructed, is of natural shape and symmetrical form, as shown on the cadaver and represented by Fig. 4. It consists of natural lip border, and is lined throughout with mucous membrane. Orbicular muscular fibres being retained in the new structure, the natural action of the lips is in a good degree preserved.

A Second Method.— Another method of reconstructing the mouth, after the loss of one-half of the upper lip and an adjacent portion of the cheek, is as follows: It consists essentially in supplying the deficiency of the upper lip by material taken from the



FIG. 5.

under lip. In such cases the under lip has itself become considerably lengthened transversely. The following are the steps of this operation: the extremity of the under lip, where it joins the right cheek, is

to be divided through its entire thickness at right angles to its border, and the division carried to the extent of one inch from the border, *a* to *b* (Fig. 5). From the terminus of this first incision, a second incision is to be extended on a line parallel to the lip border, a distance of one inch and a half toward the chin, *b* to *c*. The quadrilateral flap thus formed from the under lip is to be folded edgewise upon itself, and made to meet the remaining half of the upper lip, and be adjusted to it by its free extremity. In order, however, to make this fold, the under-lip flap must first be divided obliquely half across its base, where it still retains its connection with the chin by an incision in the line *c* to *d*.

The left half of the upper lip is also to be prepared for the new adjustment, first, by liberating it so that it can be glided toward the right side. This is accomplished by incising the buccal mucous membrane along the line of its reflection from the jaw to the lip and cheek, and detaching the parts above toward the orbit from the underlying periosteum. Second, by paring a strip of vermilion border from the extremity of the half-lip of sufficient length to permit the end of the half-lip to be matched to the free extremity of the under-lip flap. The parts concerned having been thus prepared, the under-lip flap is next to be doubled edgewise upon itself, and its free extremity adjusted to the half of the upper lip,

and the two secured to each other in a vertical line below the *columna nasi* by sutures. The space between the newly-adjusted half of the mouth and the neighboring cheek is to be closed by approximating the opposite parts and securing them to each other by sutures after their edges have been carefully matched. When the process of healing has been completed, and the parts have regained their natural pliability, the newly-constructed half of the mouth



FIG. 6.

—which has assumed a circular form and pouting shape, as shown by Fig. 6—may be restored to its natural angular shape and dimensions by a supplementary operation, which may be executed as follows: An incision is to be made with great exactness along the line of the vermilion border circumscribing the circular half of the mouth, and extend-

ing to an equal distance on the upper and lower lips (*a* to *b*). This incision should only divide the skin, without involving the mucous membrane. A sharp-pointed, double-edged knife should then be inserted at the middle of this curved incision, and directed flatwise toward the cheek, between the skin and mucous membrane, so as to separate them from each other as far as the new angle of the mouth requires to be extended. The skin alone is next to be divided with strong scissors, on a line with the commissure of the mouth outward toward the cheek (*d* to *c*). The underlying mucous membrane is also to be divided on a line opposite to, but not so far outward as, the incision made through the skin. The angle at the terminus of the incision of the mucous membrane is then to be accurately secured to the angle at the terminus of the incision of the skin, by a single thread suture. The fresh-cut edges of skin and mucous membrane above and below, that are to form the new lip borders, are to be shaped by paring first the skin and then the mucous membrane in such a manner that the latter shall overlap the former, after they have been secured together by fine-thread sutures inserted close together. By this procedure the natural shape of the mouth will be restored, as is shown in Cases I. and II. The same method is applicable where it is intended to extend the naturally-shaped mouth beyond its existing limits.

To avoid repetition, it may here be stated that the inhalation of ether was employed in all the operations reported in the subsequent pages.

The cases illustrating the subject of this volume are arranged under three classes.

First Class.—Loss of parts involving the face, and resulting from destructive disease or injury.

Second Class.—Congenital defects from excess or arrest of development.

Third Class.—Cicatricial contractions following burns.

FIRST CLASS.

LOSS OF PARTS INVOLVING THE FACE, AND RESULTING
FROM DESTRUCTIVE DISEASE OR INJURY.

CASE I.—*Reconstruction of the Mouth and Repair of the Nose after the Loss of the Right Half of the Upper Lip, the adjacent Portion of the Cheek and Ala Nasi, together with the entire Right Superior Maxillary Bone.*

CARLTON BURGAN, aged twenty, native of Maryland, and a private soldier in Company B, Permall Legion, Maryland Volunteers. The following particulars of his antecedent history were furnished by Robert F. Weir, M. D., Assistant Surgeon United States Army, in charge of the Army General Hospital at Frederick, Maryland, where Burgan had been a patient before coming to New York. He was taken sick June 5, 1862, with rheumatic pains from exposure to wet and cold while serving with his regiment. He continued ailing till July 4th, when he was sent to hospital and reported sick with typhoid fever. On the 3d of August following he

was transferred to the General Hospital at Frederick. About August 10th, although his general condition appeared to be improving, a small black slough, attended with fœtor, showed itself upon the gum at the root of the first upper bicuspid tooth on the right side; the slough spread rapidly outward toward the cheek, and inward upon the roof of the mouth. Both bicuspids and the canine tooth dropped out. The outer surface of the cheek became swollen, red, and glistening; the right eyelids swelled and closed. The gangrene continued to spread until it had destroyed the right half of the upper lip, the adjacent portion of the cheek and ala nasi; it also denuded the entire superior maxillary bone of the same side. It was ascertained that, before coming under Dr. Weir's charge, the patient had taken within the space of two weeks, for the relief of tenderness of the right side, hydrarg. massæ, gr. lxx; calomel, ℥ij; hydrarg. cum creta, ℥j. During the separation of the sloughs the fœtor was excessively offensive. After they had all come away, healthy action was established in the parts from which the sloughs had separated, and patient's general health steadily improved. On the 1st of October the right superior maxillary bone having separated was removed entire, together with the vertical plate of the os palati attached to it, and a narrow strip of bone belonging to the left maxilla adjacent to the suture

in the median line where the two maxillæ are articulated with each other. (*See Fig. 7.*) The bone itself is deposited in the Army Medical Museum, at Washington, No. 557. By the middle of October the healing parts had so far contracted that the orifice of the cavity in the face had diminished to nearly one-half its original size upon the surface. December 31, 1862, Burgan was admitted into the New



FIG. 7.

York Hospital, in a good state of health, with his face in the following condition: The right eye was sunken from atrophy of the eyeball, and the lids closed. The right half of the upper lip, the right ala nasi, and adjacent portion of the right cheek, together with the entire right superior maxillary bone, were gone, leaving an extensive opening into the

cavity of the mouth and the right nasal fossa. The margin of the cavity at the surface was formed below by the under lip, lengthened and extending obliquely upward on the right side to the middle of the cheek, where it terminated and adhered to the malar bone. From this point the margin of the cavity continued in a curved line below the inferior border of the orbit to a point on the right side of the nose, one finger's breadth below the inner canthus of the eye, and thence downward to the apex of the nose along the right side of the ridge. The columna nasi being destroyed, the lower border of the left ala and the rounded margin of the left half of the upper lip bounded the opening in this direction. Only one-third of the under lip was situated to the left of the median line, while two-thirds were situated to the right of it. The skin at the margin of the opening overlapped it, and dipped somewhat into its cavity. The inner wall of the cavity, toward the median plane, was formed by the septum nasi, deflected somewhat to the left side. The septum itself being deficient below from the absence of its cartilaginous portion, left the anterior extremity of the left inferior turbinated bone and the lower orifice of the nasal duct exposed to view. The roof of the cavity was occupied by the inferior scrolled edge of the middle turbinated bone belonging to the right nasal fossa. The outer wall of the cavity presented

a smooth surface, continuous below with the inner surface of the cheek. The tongue occupied the floor of the cavity, and was exposed to view. The palatine process of the os palati, which forms the posterior edge of the bony roof of the mouth and supports the velum, remained *in situ*, with its anterior



FIG. 8.

cicatrizated edge stretching horizontally across the middle of the cavity. The velum itself, thus supported *in situ*, performed its functions in deglutition. The teeth belonging to the left upper maxilla, except the middle incisor and one molar, were in place. Articulation was very defective, and resembled that of a person with a bad cleft palate. (*See Fig. 8.*)

In devising a plan for the repair of this extensive loss of parts, it was judged indispensable, as a pre-requisite step to any surgical operation, that some artificial substitute should be adapted to the cavity of the mouth that would supply the place of the lost maxillary bone, and afford a solid support to the soft parts that would have to be transposed for the reconstruction of the mouth and the closure of the cheek and nostrils. Mr. Thomas B. Gunning, a skill-




FIG. 9.

ful dentist of this city, to whom the case was submitted, generously undertook the execution of this delicate and difficult piece of work. The fixtures which he adapted were made of vulcanite, and consisted of two principal pieces, superposed, when in place, one above the other. The upper, or nose-piece, occupied the nasal fossa, and filled out the right half of the nose. It was hollow, and open in front and behind for the free passage of air. (Fig. 9.) The lower, or palate-piece, occupied the roof of the

mouth, and supported upon its upper surface the nose-piece. It consisted of a plate stretching across the roof of the mouth and supplying the dental arch at its margin on the right side, together with the teeth belonging to it. Its left margin took support from the existing teeth of that side, some of which it embraced. (Fig. 10.) The surfaces of both pieces, where they came in contact with the walls of the nasal and buccal cavities, were channeled with fur-




FIG. 10.

rowed lines to facilitate the flow of the secretions back into the fauces. Their accurate adjustment to each other, and to the cavities they were adapted to occupy, permitted them to be worn without causing any irritation. The light and indestructible nature of the material of which the pieces were constructed

adapted them admirably to their present use. On trial patient found he could wear them constantly with comfort, and could remove and replace them at pleasure. The improvement of his articulation and the increased facility of mastication they effected when in place afforded him the highest satisfaction.

The requisite preparations being now completed, it was decided to perform the first operation on the 26th of March, 1862. In order to guard by every possible precaution against erysipelas, and other preventable complications, the patient was placed in an outbuilding on the hospital premises which had not been in use for several weeks, and where he and his attendants would be the sole occupants.

First Operation.—Was performed as follows. The First Step of the operation was to prepare the left half of the upper lip. The lip being held on the stretch, the buccal mucous membrane was incised along the line of its reflection from the upper jaw to the lip and cheek as far outward as the molar teeth. The lip itself was then divided through its entire thickness from the point where it joined the left ala nasi, on a line parallel with the lip border, outward to the middle of the cheek. The lip flap thus formed was trimmed square at its free extremity. The Second Step was to prepare the redundant under lip, so as to employ it for supplying the deficient right half of the upper lip. This was done

according to the method described on page 26. After the newly-adjusted parts had been secured by the requisite sutures, the reconstructed right half of the mouth assumed a circular shape, and stood prominently forward. The Third Step had for its object to close the open space in the right cheek, resulting from the transposition of the parts effected by the Second Step of the operation. It was executed as follows: A transverse incision was carried through the entire thickness of the right cheek on a line with the commissure of the mouth, as far outward as the anterior edge of the masseter muscle, and beyond it only through the skin covering the muscle. In order to glide forward the divided cheek, the buccal mucous membrane alone was divided along the anterior edge of the masseter, above and below the transverse incision. This allowed the cheek to come forward and meet the outer margin of the transposed under-lip flap. Their confronting edges, after being pared and matched to each other, were secured in exact coaptation by silver-wire and fine-thread sutures, inserted close together. Before closing the parts, the numerous ligatures that had been applied to bleeding vessels were loosely twisted into a skein, and brought out to the surface at the outer angle of the transverse incision. The operation now completed had occupied nearly three hours, owing in part to the frequent interruptions necessary to maintain the effect of the

ether. No adhesive plaster was applied. Warm-water dressings were directed to be kept to the parts.

March 23d.—Progress favorable; pulse 90. Inflammatory tumefaction moderate. The yarn upon the pins in the upper lip being sunken in the swollen parts, it was removed, and, after allowing the underlying constricted surface to recover itself, fresh yarn was applied. Liquid nourishment and drinks were taken without difficulty. Patient experiences no uneasiness from the presence of the artificial fixtures. Injections were employed to cleanse the spaces between and around them.

28th.—General condition favorable; pulse 90. Swelling extended to the right side of the neck, below the jaw. No indications of deficient vitality existed at any point. Removed the alternate thread sutures and the silver-wire sutures from the lip.

29th.—Progress still favorable; pulse 104. Copious suppuration is taking place from the track of the ligatures, and is somewhat offensive. Removed additional thread sutures. At the junction of the two halves of the upper lip the new adhesions were in danger of giving way, owing to tension. To prevent it a fresh pin was inserted between the two old ones, which were then removed.

30th.—Several ligatures have come away. Suppuration diminishing and no longer offensive; removed the sutures from the vermilion border.

31st.—Removed the supplementary pin suture, and supported the upper lip with adhesive plasters.

April 9th.—Parts had all healed, and dressings were no longer required. The result is shown by Fig. 11.



FIG. 11.

The parts involved in the preceding operation having regained their natural suppleness, a Second Operation was performed April 23d, for the purpose of restoring to the right half of the mouth its natural angular shape, and lengthening it at the same time.

Second Operation.—Was performed according to the method described on page 28. At the expiration of twenty-four hours after the operation the alternate

thread sutures were removed, and on each succeeding day others were got rid of, as fast as they could be dispensed with, until the fourth day, when healing was complete, and the last suture was removed. The result was satisfactory, and is shown by Fig. 12.




FIG. 12.

The dental fixtures having been worn constantly for more than six weeks, it was found necessary to remove them, in order to relieve the natural teeth of the left upper maxilla from the too great pressure to which they had been subjected by the insertion of pieces of sponge between the right cheek and the artificial teeth belonging to the palate-plate; these sponges had been employed to resist the contraction

of the newly-cicatrizing parts. During the succeeding eight weeks, in which the dental fixtures were left out of use, the parts underwent changes which require some particular notice. For instance, wherever the incisions involving the right cheek had divided the lining mucous membrane, tense salient bands had formed. One of these crossed the inside of the cheek horizontally, another extended upward and backward, deep below the orbit; both bands were continuous in front with the upper border of the transposed under-lip flap, which now constituted the right half of the upper lip. This same border also formed the lower margin of the still-existing opening into the right cheek, which corresponded to the deficient right ala nasi. In attempting to open the mouth, these bands were put tightly on the stretch, and, proceeding from a point above the outer incisor tooth of the left upper maxilla, they prevented a separation of the jaws beyond a space sufficient to admit a finger edgewise between the teeth. A firm adhesion between the cheek and lower jaw, on the right side of the chin, was an additional obstacle to the separation of the jaws. This condition of the parts was an insuperable obstacle to the re-introduction of the dental fixtures, and required to be overcome preparatory to any further operation. This was done without anæsthesia in the following manner: A pair of blunt-pointed scissors, curved

flatwise, was introduced into the mouth, guided by the forefinger of the left hand, and with them the upper band was divided at its remotest point under the orbit. The horizontal band was also freely divided at its farthest point, and the adhesions between the cheek and lower jaw were also freely liberated. Considerable relaxation of the parts was thus obtained, but not sufficient for our purpose, until the left half of the upper lip was a second time divided through its entire thickness horizontally along the line of the cicatrice left by the first operation. This had the desired effect, and the replacement of the dental fixtures was now accomplished by Mr. Gunning himself, who was present on the occasion.

Third Operation.—Performed June 18th, at 3 P. M. The object of it was to close the remaining opening in the right cheek, and cover the adjacent side of the nose with a patch of skin, to be taken from the left half of the forehead. The opening, as shown by Fig. 12, involved the right half of the nose below the os nasi, and the adjacent cheek as far out as its middle. The lower border of the opening extended in a line from the inferior edge of the left ala nasi horizontally across to the middle of the right cheek, and formed also the upper margin of the newly-transposed under-lip patch. In order better to adjust the edges of the patch of skin that was to be brought down from the forehead, the skin at the

margin of the opening was dissected up from its underlying connections sufficiently to permit it to be everted. At the outer margin of the opening, which corresponded to the anterior edge of the masseter muscle, the skin, which dipped somewhat into the cavity, was detached more extensively, so as to permit it to be glided forward, and thus contribute to the closure of the opening. A pattern of the shape of the opening, but somewhat exceeding it in size, as an allowance for shrinkage, was then cut from oiled silk and laid upon the left half of the forehead, in an inverted position, with its broadest part directed upward, and its narrowest applied above the inner half of the left eyebrow. An incision was then commenced at the tip of the nose, Fig. 12, *a*, and carried upward along its dorsum, skirting the margin of the opening, and extending beyond it obliquely to the inner extremity of the left eyebrow, *b*, where it encountered the right margin of the pattern, and was continued thence around its entire circumference till it reached the middle of the left eyebrow, *c*. The included patch of skin was then dissected up from the pericranium, and left attached below at the supra-orbital ridge by its pedicle, which measured one inch in breadth. In order that the patch from the forehead should, after its transfer, have its under surface lie in immediate contact with raw surface, the skin intervening between the eyebrows was dissected up

and removed. The patch itself was then brought round edgewise from left to right, and above downward, till it reached its new location on the right side of the nose and cheek, where it was adjusted with its straight edge along the entire length of the dorsum nasi, and secured there by three pin sutures, inserted at equal distances apart, and additional intermediate fine-thread sutures. What had now become the right border of the patch in its new locality, was adjusted to that part of the margin of the opening which involved the cheek, and was secured to it by fine-thread sutures, inserted close together. The inferior and outer angle of the patch, which, after its junction with the cheek, would bound the nasal orifice on the right side, was adjusted as follows: At the angle where the outer and inner borders of the opening in the cheek met, a strip, half an inch long, was raised from the inferior border, and left attached toward the median line. The angle of the patch was then adjusted in its place, and the detached strip applied to its inner raw surface, and made to line it. The edges of the transverse incision, by which the left half of the upper lip had been liberated the second time, were again secured to each other by sutures. The ligatures that had been applied to bleeding vessels were brought to the surface at the nearest point of exit. The raw surface left upon the forehead was coated with a col-

lotion crust, in the manner described on page 13. Lint was stuffed between the cheek and jaw, on the right side of the chin, to prevent adhesion from again taking place. Warm-water dressings were applied to the forehead and cheek, and dry warmth to the nose, by means of a bat of cotton and a vial of hot water suspended in contact with it. Although the operation occupied nearly three hours, it was not followed by any unpleasant effects from the long-continued administration of ether.

June 19th.—Patient passed a comfortable night, and obtained some sleep. The transferred patch was cool to the touch, the difference of temperature being 10° . Covered the surface with threads moistened with spirits of turpentine, and continued the same applications.

20th.—Progress favorable; temperature of the patch was restored and well sustained. The confronted edges of the patch and adjacent skin were alike tumefied, and showed no difference in their degree of vitality. Changed the yarn on the pins.

22d.—Patient continued to do well. Removed the only remaining pin from the tip of the nose, and the alternate thread sutures at other points.

23d.—Removed other sutures; all the ligatures had come away.

24th.—The edges of the patch had united at all points by first intention.

26th.—The last suture was removed. Strips of adhesive plaster were applied to support the newly-healed parts.

27th.—This being the ninth day, the collodion crust separated and dropped off, exposing a healthy granulating surface, nearly on a level with the surrounding skin. Scarcely any inflammatory redness or swelling had at any time appeared around the margin of the crust.

July 24th.—Continued favorable progress. The transferred patch in its new connections had undergone a process of hypertrophy, and obtained a uniform thickness of half an inch. The adjacent skin, to which it was united, participated in the same thickened condition to a distance of one-quarter of an inch from the line of junction. The surface of the patch was paler than the neighboring skin; the sore upon the forehead had progressively diminished in size. Upon the upper part of the nose, and between the eyebrows, the skin was gathered into a bulging fold, in consequence of the doubling of the pedicle of the patch upon itself in its transfer from the forehead. The conspicuous disfigurement thereby produced was now to be removed by a

Fourth Operation.—Performed August 8th. Two parallel incisions, commencing one on either side of the dorsum nasi, were continued upward on the forehead, where they converged and met so as to in-

clude the bulging fold of skin in a tongue-shaped strip, which was itself dissected up from its underlying connections. Narrow strips of skin were removed from the edges of the wound on either side, to make room for the strip, which was spread out and replaced in an advanced position, and secured by numerous sutures to the adjacent skin on all sides. With a view to reduce the excessive thickness of the new ala nasi, at its inferior margin, where it bounded the orifice of the nostril, a prism-shaped strip, penetrating deep between the outer and inner surfaces, was excised, and the wound closed by fine-thread sutures. Both these operations were followed by primary union, and did well.

September 1st.—The wound surface upon the forehead had healed, leaving a cicatrice scarcely exceeding one-third the size of the original wound. The shrinkage which the nose-patch had undergone had produced a deep furrow along the cicatricial line, occupying the lower third of the ridge of the nose. The furrow terminated below, at the apex of the nose, in a deep, unsightly notch. To remedy this disfigurement was the design of a fifth, and final, operation.

Fifth Operation.—Performed October 27th, as follows: Two parallel incisions were made, one on either side of the furrow, penetrating deep in converging planes, so as to include both sides of the furrow and the notch below. The opposite edges of the wound

were secured in accurate contact by two pin sutures and three fine-thread sutures. Within six days the sutures were all removed and perfect union obtained. A great improvement in the appearance of the nose was the result of this operation. In June, 1864, Bur-
gan was in the enjoyment of good health, and had for several months preceding discharged the duties of an assistant-nurse in a large ward of the hospital. The hypertrophied condition of the nasal patch still persisted giving to the right half of the nose a plump form. When the surface was pricked, the sensation was no longer referred to the forehead as at first, but to the actual seat of irritation. The cicatricial bands on the inside of the right cheek had been kept from again contracting by the persevering efforts of the patient, who had faithfully executed the directions given him, to introduce one or more fingers into the mouth and stretch the bands to their utmost limit, and to repeat the operation several times daily. The artificial substitute for the right maxilla, which consists of a palate-plate that covers the roof of the mouth, and supplies the lost teeth of the right side, is the only piece now worn, the nasal piece having been dispensed with for a long time. It is worn constantly with entire comfort, and can be removed and replaced at pleasure. With it patient is able to masticate all kinds of food, and articulation, which without it was scarcely intelligible,

now betrays very little defect. In May of 1871 the author visited Burgan at his home, in the suburbs of Baltimore, and found him a married man, and father of two children. He enjoyed good health, and was pursuing a laborious occupation. He still wore constantly the same palate-plate that had been



FIG. 13.

adapted in March, 1863. Time had considerably improved the condition of the right cheek and nose. The cicatricial lines on the surface of the cheek were no longer elevated ridges, but had shrunk into shallow furrows, and become much less conspicuous. The cheek itself was pliable at all points, and the right half of the nose retained its plump shape, al-

though it had lost its excessive thickness. Fig. 13 is from a photograph, taken at Balch's Studio in Baltimore, in 1871.

CASE II.—*Reconstruction of the Mouth and Repair of the Nose after the Loss of the Right Half of the Upper Lip, the Adjacent Portion of the Cheek, and Right Ala Nasi.*

JOHN MICHÆLIS, aged eleven, of German parentage, a resident of Jamaica, L. I., was admitted into St. Luke's Hospital in April, 1866. The loss of parts sustained in this case happened in the progress of malarial fever, with which he was attacked in the month of September preceding. After a fortnight's treatment, during which, according to his mother's statement, he took a great deal of medicine, sloughing invaded the right half of the upper lip and adjacent cheek. At the expiration of about ten weeks the parts from which the sloughs had separated were healed, and had assumed the condition in which they were at the time of his admission to the hospital. It was as follows: The right half of the upper lip, the adjacent portion of the right cheek, and the right ala nasi, were gone. The upper front teeth and gum-surface belonging to the right half of the upper maxillary bone were bare and exposed to view. A

supernumerary canine tooth emerged from the gum above the level of the other teeth, in the space between the outer incisor and canine teeth, and formed a conspicuous feature in the disfigurement of the face. The gum-surface above these teeth was incrustated with a brownish scab, from underneath which healthy pus escaped on pressure. On the right side of the nose a semicircular notch occupied the place of the ala that had been destroyed. The upper edge of the notch corresponded to the lower margin of the os nasi. The septum and columna nasi were entire. The cicatrized margin bounding the deficiency in the cheek was sunk, and closely adherent to the upper maxilla above the teeth. The lower-lip border was lengthened on the right side, and stretched obliquely outward and upward to the alveolar border of the upper jaw, where it adhered to the gum-surface above the bicuspid teeth. From this point of adhesion a linear cicatrix one inch long extended outward, and a little downward, half across the cheek, depressing the surface and adhering to the underlying parts. The median line divided the under-lip border unequally, three-fifths of the border lying to the right and two-fifths to the left of it. The remaining left half of the upper lip was shrunk, at the point where its vermilion border terminated below the columna nasi. The motions of the lower jaw were unrestricted. The patient's general health

was good. The outer incisor, canine, and supernumerary canine teeth of the upper jaw were extracted preparatory to an operation. Fig. 14 shows the condition of the face just described.




FIG. 14.

First Operation was performed May 18th as follows: The left half of the upper lip, being held upon the stretch, was detached from the jaw by an incision of the buccal mucous membrane, carried along the line of its reflection from the jaw to the lip and cheek, and extended outward as far as the molar teeth, and also upward on the level of the periosteum toward the orbit. This permitted the lip and cheek

to be glided over toward the right side. A strip of the vermilion border, one inch in length, was pared away from the extremity of the half lip and left attached temporarily to it. Material with which to supply the deficient half of the upper lip was next obtained from the redundant right half of the under lip. It was done according to the method described on page 26, and also employed in Case I. (*see* page 38). The quadrilateral flap (*a, b, c*) thus formed from the under lip was adjusted by its free extremity, after being brought round edgewise to the left half of the upper lip, and the two were secured together in a vertical line below the *columna nasi* by two pin sutures and several fine-thread sutures. The open space in the right cheek, remaining after the transposition of the parts as just described, was now closed in the following manner: An incision, one inch and a half long (*e to f*), was carried transversely across the right cheek, on a level with the middle of the nose, to a point one inch below the outer canthus of the eye, and thence downward in a curved line (*f to g*), with its convexity directed backward to a point within one inch of the margin of the lower jaw. The quadrilateral flap of skin (*b, e, f, g*), lined with mucous membrane, was dissected up from the jaw, but left attached below. It was then slid forward edgewise till its anterior border met the outer border of the newly-transposed under-lip flap, and

the two were matched together and secured by pin sutures and fine-thread sutures, distributed so as to effect the most exact adjustment of the edges to each other. In order to cover the bare surface remaining on the cheek after the transfer just made, the incision *e, f*, crossing the cheek, was prolonged to the point *h*, upon the temple, and the angle of skin, *h, f, g*, included between it and the vertical incision, *f, g*, was dissected up from its underlying connections sufficiently to allow it to be drawn forward and cover the bare surface, and there be adjusted with sutures. The reconstructed right half of the mouth had now assumed a circular and pouting form, a defect which it was intended to remedy by a subsequent operation. The same after-treatment was employed in this case as in the preceding one, and with a like good result. There was no lack of vitality, nor sloughing, at any point. The inflammatory tumefaction was moderate, and the sutures were removed in succession as they could be dispensed with. On the tenth day all had been got rid of, and patient was out of bed and able to go about. The result of the operation is shown in Fig. 15. The notch involving the right ala of the nose, constituting as it did a part of the original defect, now formed with the adjacent cheek and upper lip a foramen capable of admitting the end of a finger. The closure of this foramen was next to be

attempted by a second operation, which was performed on the 18th of June, as follows :




FIG. 15.

Second Operation.—The edge of the foramen, where it involved the cheek and upper lip, was pared and everted. On the right side of the apex nasi an incision, skirting the edge of the notch, was carried up obliquely over the dorsum of the nose to the inner extremity of the left eyebrow, *a* to *b*. A second incision, commencing at the opposite edge of the foramen upon the right cheek, was carried upward upon the right side of the nose parallel with the first incision to the inner extremity of the right eyebrow. The skin included between these two in-

cisions was dissected up from the side of the nose, and between the eyebrows, and removed.¹ A pattern of the shape of the bare surface thus prepared was cut from oiled-silk, and laid upon the forehead above the inner half of the left eyebrow. The incision previously made, that terminated at the inner extremity of the left eyebrow, was then continued upward along the right edge of the pattern, and onward around the rest of its margin to the middle of the eyebrow. The included patch of skin was then dissected up from the pericranium, but left attached below at the margin of the orbit where its pedicle derived a vascular support from a branch of the ophthalmic artery emerging from the orbit. The patch of skin was now brought down edgewise from left to right to its destined locality, and its edges accurately adjusted to the edges of the space prepared for it, by pin sutures and fine-thread sutures inserted close to each other. A short piece of gum-catheter was inserted in the nasal orifice, and maintained there for the purpose of establishing a permanent aperture. The opposite edges of the bare surface upon the forehead were approximated, and held in contact by sutures and strips of adhesive plaster.

¹ This portion of skin it would have been better to save and utilize for covering the bare surface on the forehead, as was done with great advantage in subsequent similar operations.

The after-treatment was the same as already described in previous operations. The subsequent progress was favorable. The transposed patch maintained its vitality, and united by primary adhesion at all points except at its lower extremity, where, after suppurating for several weeks, it healed at



FIG. 16.

length by the granulating process. The wound upon the forehead failed to heal by adhesion, but finally healed by granulation, the edges being held approximated during the process by adhesive-plaster, so that the resulting cicatrice was reduced to almost linear dimensions. The result of the operation is shown by Fig. 16.

On the 31st of July, patient was allowed to go to his home to recruit. After his return to the hospital much improved in health, a third operation was undertaken for the purpose of restoring the newly-reconstructed right half of the mouth, which had assumed a circular, protruding form, to its natural angular shape, and at the same time to lengthen the mouth on the same side.

Third Operation was performed September 26th, according to the method described on page 28. An operation was also performed at the same time for the removal of a conspicuous ridge of skin occupying the upper part of the nose, and the space above between the eyebrows. It had resulted from the folding upon itself of the pedicle of the patch of skin that had been brought down from the forehead. Two curved incisions were carried, one on either side of the elongated ridge, and made to meet at its opposite extremities. The elevated skin included between the incisions was removed, and the opposite edges of the wound were brought together and secured by sutures. Primary adhesion followed in both operations, and an excellent result was obtained.

A great improvement of the external appearance of the right half of the face had been effected by the operations already performed, but there still remained a serious defect dependent on cicatricial con-

tractions on the inside of the right cheek, which held the jaws together and prevented the teeth from being separated in front more than half an inch. A single cicatricial band of mucous membrane on the inside of the right cheek, close to the angle of the mouth, presented the greatest resistance to the separation of the jaws. With the view of remedying this defect it was proposed to lengthen the mouth, which was scanty on this side, by extending the angle farther toward the cheek. In doing this the constricting cicatricial band on the inside of the cheek would be divided, and the jaws thereby liberated. For this purpose a fourth, and final operation, was performed in July, 1868.

Fourth Operation was a repetition of the preceding third operation, and secured the desired result, so that the mouth became more symmetrical in shape, and the teeth could be separated one inch in front.

Before leaving the hospital the patient was instructed to insert wedges of wood between the upper and lower molar teeth, and wear them as long at a time, and as often daily, as possible, with the view of increasing the power of separating the jaws. A progressive improvement was afterward observable. In April, 1871, the mouth could be opened and the jaws separated to the fullest extent. The right cheek was pliable, the cicatricial lines on its outer surface



FIG. 17.

had shrunk and become much less conspicuous. Fig. 17, copied from a photograph taken at this time, shows the final result.

CASE III.—*Reconstruction of the Nose and Mouth after the Loss of the Nose and entire Upper Lip.*

JANE TUCKER, aged twenty-six, native of Ireland, unmarried, admitted into St. Luke's Hospital, March 10, 1866. The antecedent history of her case is as follows: When about seven years old, sores formed on the inside of her nose, from a practice she was

addicted to of constantly picking her nose with her fingers. Destructive ulceration followed, which extended to the upper lip, and finally resulted in the condition in which she was when admitted into the hospital. It was as follows: The nose below the nasal bones was sunk to the level of the cheeks, and both nostrils were blocked up, the only external opening being a small perforation situated in the median line on a level with the floor of the nasal cavity, and of a size capable of admitting an ordinary probe. The ossa nasi still remained *in situ*. The upper lip was entirely gone, leaving uncovered the upper front teeth and adjacent gum-surface within an angular space bounded by two lines which diverged from a central point on a level with the floor of the nostrils, and descended to the angles of the mouth. These lines also marked the limits of the sound skin on either side. Both angles of the mouth were entire, and the under lip was of ample dimensions, being even somewhat pendulous. Patient's general health was good. Fig. 18 shows the condition just described.

First Operation.—The design of the operation was to reconstruct the upper lip from the redundant material of the lower lip. It was performed on the 28th of March, as follows: An incision, commencing at the median line, on a level with the floor of the nasal cavity, was carried outward and downward on

both sides of the face in a curved line so as to circumscribe both angles of the mouth, and terminate at a point below the junction of the middle and outer third of the under lip (*a* to *b*, *a* to *c*, Fig. 18). These incisions divided the entire thickness of the cheeks and lip, and in their course were kept at a



FIG. 18.

uniform distance of one inch and a quarter from the angles of the mouth and the under-lip border. The flaps thus formed on either side were brought toward each other edgewise, and their ends, after being pared and made straight, were adjusted to each other on a vertical line in the median plane, and secured

by three pin sutures and intermediate thread sutures. What had now become the upper border of these united flaps was adjusted to the opposite cut edge of the skin above, and secured by sutures. In order to close the open spaces between the newly-transposed flaps and the cheeks on either side, from which they had been detached, it was necessary to liberate the cheeks by incising the mucous membrane on the inside of the mouth, above and below, along the line of junction between the cheeks and the jaw-bones. This permitted the cheeks to be brought forward to meet the flaps encircling the new mouth, and close up the spaces between them. This adjustment was secured by pin sutures and thread sutures distributed so as to afford support to the parts at all points, and effect the most exact coaptation of their edges. No bare surface was left uncovered. The reconstructed mouth necessarily assumed a circular and pouting shape. Primary union followed at all points, except where the flaps came together in the median line below the nasal cavity. Here it failed in consequence of sloughing of the newly-united edges, which separated from each other except at the upper third of their line of junction. This separation, including as it did the lip border, left the upper front teeth again uncovered as before, though not to so great an extent.

Patient remained in the hospital till the 24th of

July, when she was discharged in ordinary good health. On the 13th of March, 1867, she was admitted into the New York Hospital to undergo further operations. It was now proposed to remove the obstruction of the nostrils, and afford a free passage for air through them, thereby enabling the patient to have an artificial nose adapted.

Second Operation.—A vertical incision was carried from a point midway between the eyebrows downward upon the nose to a point on a level with the floor of the nasal cavity; a transverse incision crossed the lower extremity of this vertical incision, and extended one inch on either side of it, the two forming together an inverted letter-T-shaped incision. The angular flaps of skin on either side were dissected up, and the sunken parts blocking up the nasal orifice were cleared away from between the ascending nasal processes of the superior maxillary bones, which form on either side the lateral boundary of the nasal fossæ. The skin was then pared to correspond to the bony margin of the new opening and left to cicatrize, which it did in due time, leaving a permanent aperture with rounded edges, and of the shape of an inverted heart, measuring three-quarters of an inch in its vertical, and five-eighths of an inch in its transverse, diameter. (*See Fig. 19.*)

Third Operation was performed on the 14th of

April, 1867. Its object was to attempt once more the reconstruction of the upper lip. Although the first operation on the mouth had been only partially successful, an important advantage had been gained by securing a covering of skin between the upper front teeth and the nasal orifice. The steps of the operation were the following: An incision was commenced about half a finger's breadth below the nasal orifice and carried along the margin of the newly-transposed skin, and continued on both sides outward and downward in a curved direction, so as to circumscribe the angles of the mouth, and terminate at corresponding points on each side of the chin, below the junction of the middle and outer third of the under-lip border. These incisions were kept at a distance of one inch from the angles of the mouth and the under-lip border, and divided the entire thickness of the cheek. The two lateral flaps thus formed were brought together edgewise and made to meet in the median line, where their ends, after being squared, were adjusted together by sutures. The outer (i. e., toward the cheeks) edges of these united flaps were readjusted to the opposite edges of the cheeks, from which they had just been detached, by the same procedure as had been employed in the first operation, of which this was almost an exact repetition. As a result of this operation the mouth again assumed a circular and pouting shape.

The subsequent management of the sutures was the same as has been already described in previous operations. The edges of the flaps at their junction in the median line below the nasal aperture, and at their junction with the cheek, to the distance of one inch on either side of the median line, assumed a livid, ashy appearance, and failed to unite by primary adhesion.

Erysipelas of a mild form showed itself on the 16th of April, attended with moderate swelling; on the 19th it had disappeared. The sloughy edges of the wound having cleared off and begun to granulate, it now became imperative to maintain them in contact during the process of union by granulation. This was effected by sutures, inserted at every point where they could afford support. General invigorating treatment was also enforced, such as sulph. quiniæ, iron, wine, and generous diet.

From this date onward to the 7th of May, the most assiduous care was required to support the edges in contact with each other. Sutures were renewed at every available point once in two or three days, and removed before they had time to produce suppuration. Strips of adhesive-plaster were also employed for support. Union was at length secured at all points. The left half of the new upper lip, having suffered most from sloughing, is somewhat shrunk, and also slightly notched, at its border,

where it joins the right half in the median line. The result thus achieved, though not perfect, has nevertheless brought the parts into a condition in



FIG. 19.

which it will be possible by another operation to restore the mouth to its natural shape.

Fourth Operation.—The mouth having assumed, as a result of the preceding operation, a circular form, it was the design of the fourth operation to restore it to its natural angular shape. It was executed in July, 1867, on both sides of the mouth simultaneously, according to the method described on page 28, and was followed by a satisfactory re-

sult, as shown by Fig. 19. After leaving the hospital in January, 1868, the patient had an artificial nose, made of vulcanite, adapted to her face. She wore it for some time, but without any satisfaction, and finally cast it aside in disgust.

On the 8th of January, 1869, Jane was again admitted into the New York Hospital, to undergo an operation for the construction of a nose, the material for which was to be taken from the forehead.

Fifth Operation.—As a preliminary step to this operation, a nose was modeled in wax by a sculptor upon a plaster cast of the patient's face previously taken for the purpose. This served as a pattern from which to determine the size and shape of the patch of skin that would be required. The external nasal aperture was then prepared by carrying an incision along its margin on both sides, and dissecting up the outer edges of these incisions toward the cheeks, so as to adapt them for coaptation with the lateral edges of the patch of skin that was to form the new nose. A pattern, cut from oiled-silk and shaped upon the model of the nose, was laid upon the forehead in an inverted position, with its pedicle of attachment applied above the left supraorbital notch, and its long axis lying obliquely upon the forehead and inclining to the right of the median line. An incision, involving the thickness of the skin, was then carried around the margin of the pat-

tern, and the included patch of skin, after being dissected up from the pericranium, was left attached below at the margin of the orbit. The skin covering the space between the eyebrows and the ossa nasi was displaced, to afford a continuous raw surface to which to apply the under surface of the patch, when brought down from the forehead. This portion of skin, however, was left attached at the inner extremity of the right eyebrow and reserved for subsequent use. The patch itself was then brought down edgewise from the forehead to its destined location over the nasal aperture, to the edges of which (prepared as described above) the edges of the patch were accurately adjusted and secured by numerous sutures. The bare surface left upon the forehead was covered at its lowest part by the reserved portion of skin that had been displaced from between the eyebrows. The upper portion of the bare surface was treated by covering it with a collodion crust (*see* page 13). The cavity of the newly-constructed nose was stuffed with lint to maintain it plump and in good shape. The patch, in its new locality, underwent no apparent change in color or temperature. Primary union followed at all points except on the right side of the nose, high up, where suppuration took place between the edges of the wound, for a distance of three-quarters of an inch.

Several weeks after the operation just described,

an attempt was made to close the opening by paring its edges and securing them in contact by sutures. To my great disappointment sloughing followed, and an enlargement of the size of the opening was the consequence. After a further delay of several months the opening still existed, and measured three-fourths of an inch in its vertical, and half an inch in its transverse, diameter. The closure of this opening was next attempted by a

Sixth Operation.—It was executed as follows: The edges of the opening were pared and everted. A patch of skin, of suitable shape and size, was raised from the forehead above the opening, and left attached below at the supraorbital notch on the right side. It was then turned down and adapted to the opening, with its cuticular surface directed toward the nasal cavity and its raw surface outward. Fine-thread sutures were inserted in close proximity to each other, to secure the most accurate adjustment of the edges. In making this transfer it was necessary to stretch the flap across an intervening space of sound skin of about a finger's breadth.

At the end of three weeks, the patch, which had become ingrafted in its new locality, was divided at the upper margin of the opening and its pedicle turned up and replaced in its original site, there to contribute toward covering the bare surface upon the forehead from which it had been taken. A com-

plete closure of the opening, however, was not effected by this operation, as a small aperture still remained at the upper margin, allowing the passage of air through it. The outer raw surface of the transplanted patch cicatrized, and adapted itself so exactly in its new locality that its outline could scarcely be distinguished.

In the month of June, 1870, Jane again became a patient at St. Luke's Hospital, and two successive operations were performed for the purpose of closing the opening still existing on the right side of the nose. The first of these operations consisted in paring the edges of the opening, detaching them extensively from the underlying parts and securing them in contact by sutures. This failed from defect in the adhesive process.

After waiting a sufficient length of time, and allowing the edges to cicatrize, a second attempt at closure was made as follows: The edges of the opening were seared with the actual cautery, and, as soon as they had assumed a granulating condition, a slender tenotomy-knife was inserted under the skin, at a distance of three-fourths of an inch from the opening, and swept around flatwise in every direction so as to liberate the skin from its underlying connections. In order to maintain the opposite granulating edges of the opening in contact, and thus facilitate their adhesion, a beaded silver-wire suture (*see* page 17)

was inserted crosswise. This expedient, although it secured the desired contact of the edges, failed to effect a complete closure of the opening. What remained of it, however, ultimately contracted down to so small a size that the patient was contented to wear a small plug of wax which kept it closed, and was scarcely noticeable. She also wore habitually a plug of lint in the nasal cavity to keep it distended and maintain the nose more prominent.



FIG. 20.

In the summer of 1871 the patient's condition was ascertained to be as follows: The mouth retained its natural shape and dimensions, and the

lips performed all their functions. The nose had shrunk considerably since it was first reconstructed, though for a year past it had remained stationary. The voice was nasal in a very marked degree. The nose, though defective in shape, was much less repulsive to the eye than an artificial nose would be. Fig. 20, showing the final condition of the face, is from a photograph, as are the other two.

The patient was again examined June 23, 1875. The condition of her face remained much the same as described above, without having undergone any perceptible deterioration.

CASE IV.—*Reconstruction of the Mouth after the Loss of the entire Under Lip, and a portion of the Inferior Maxillary Bone.*

HUGH B., aged thirteen, a resident of the city, of rather slender constitution, though ordinarily enjoying good health, came under surgical treatment in October, 1869, for the relief of the condition of his mouth, of which his mother gave the following account: When six years old he was attacked with scarlet fever, and became dangerously ill. The eruption was scanty, and in other respects the disease was irregular in its development. Sloughing of the under lip followed, accompanied by inflammation in-

volving the region of the lower jaw on both sides of the face. Necrosis supervened, and considerable portions of the jaw-bone were cast off. The condition of his face consequent thereupon was as follows: The entire under lip was gone, and the skin below it. On the right side of the chin the loss of skin extended to within one finger's breadth of the edge of the jaw, while on the left side it did not extend so low down. The upper lip was of ample dimensions, and both angles of the mouth were entire and somewhat drawn down. The right half of the tongue adhered by its under surface to the floor of the mouth, as far forward as the alveolar margin of the maxilla in front; its extremity, which was also bound down by adhesions, was exposed to view from the absence of the under lip. That portion of the body of the lower jaw between the symphysis and last molar tooth on the right side had previously come away entire, and the alveolar sockets of all the teeth belonging to it could be identified in the bony specimen (*see* Fig. 21). The last molar tooth on the right side alone remained *in situ*. A much smaller portion had also come away from the left half of the jaw. It consisted of the alveolar border supporting the sockets of the two bicuspid and two adjacent molar teeth (*see* Fig. 22). This extensive loss of bone on the right side had been supplied by new bony product along the entire inferior border of the maxil-

la to such a degree that the symmetrical form of the face was maintained and a solid support afforded. The canine tooth on the left side of the lower jaw and the last molar on the right side were the only



FIG. 21.



FIG. 22.

teeth remaining *in situ*. Articulation was but little affected. The greatest discomfort experienced by the patient was from the constant dribbling of saliva. Owing to the loss of teeth his nourishment was restricted to liquid and soft solid articles of food. His general health was good. (*See* Fig. 23, taken from a plaster cast.)

First Operation.— Was performed October 5, 1869, at patient's residence, with the aid of Prof. A. C. Post and Drs. C. M. Bell and J. N. Beekman. It was executed as follows: The right cheek was dissected up on the inside of the mouth, from the jaw downward as far as the lower border of the

jawbone, and backward to a point a little beyond the anterior edge of the masseter muscle. An incision was then carried from a point, *a*, below the middle of the zygoma downward upon the right cheek, and forward in a curved line to a point, *b*, half an inch below the right angle of the mouth. After dis-




FIG. 23.

secting up the flap thus formed, and getting access to the cavity of the mouth, the mucous membrane alone was divided along the anterior margin of the masseter muscle upward, and then forward as far as the upper canine tooth. By this procedure the entire cheek flap was liberated, and could be advanced edgewise forward till its free extremity, carrying

with it the angle of the mouth, reached the line of the symphysis menti. The same procedure was executed on the left cheek. The arteries were ligated as fast as they were encountered, thus sparing hæmorrhage. The skin covering the prominence of the chin was shaped symmetrically into an angular form, the angle pointing upward, and occupying the median line. The two cheek flaps were then glided edgewise forward, and made to meet by their anterior edges over the symphysis, where they were secured together by two pin sutures, inserted below the lip border, and three fine-thread sutures at the border. Below their line of junction over the symphysis the edges of the two cheekflaps diverged from each other, and stood astride of the angle of skin covering the chin, and were adjusted to it by sutures. The effect of this adjustment was to draw the angles of the mouth toward each other, and fold the upper lip upon itself, so that it stood forward in advance of and overhung the short, retracted under lip. The surfaces left bare on both cheeks by the transfer of the flaps, as just described, were covered by dissecting up from their underlying connections the edges of the wound from which the flaps had just been detached, and gliding them forward to meet the cheek flaps, and be again adjusted to them by sutures. By this means, at the completion of the operation, there remained no uncovered, bare surface. Notwithstand-

ing the extensive dissections required in this operation, the hæmorrhage was controlled within moderate limits, and was not followed by any marked depression of the pulse. A single attack of vomiting, which emptied the stomach of the blood swallowed during the operation, was the only disturbance produced by the protracted administration of ether. The parts were covered with a layer of double thickness of sheet-lint to maintain their natural warmth. The inflammatory tumefaction developed during the three or four days succeeding the operation was moderate, as was also the febrile reaction. On the second day I began removing the alternate thread sutures, and changing the yarn on the pins. Special care was required to maintain in close contact the edges of the flaps that were united over the symphysis and on both sides of the chin, so as to prevent, if possible, the escape of saliva between the sutures. On the right side of the chin, where the tongue came in contact with the parts on the inside of the mouth, and exerted some pressure against them, saliva did escape, and prevented adhesion from taking place. At all other points, however, primary union was secured, but not without renewing the sutures at different points, whenever ulceration began to take place, before the support of the original sutures could be dispensed with. The administration of nourishment was managed by patient's mother, whose long experience en-

abled her to do it very successfully. Union having failed to take place on the right side of the chin, there remained a narrow aperture of nearly two inches in length, with cicatrized, rounded edges, that required an operation for its closure.

Second Operation.—Performed October 30th, as follows: Both edges of the aperture were pared afresh. The lower edge was dissected up from the bone and everted; the upper edge was cut across at both ends, so as to permit it also to be everted. The two edges were then accurately confronted and secured together by six pin sutures and nine intermediate silver-wire sutures, the wound thereby being rendered impermeable to saliva.

November 4th.—The last suture was removed and union obtained at all points, except one, where an opening of the size of a goose-quill allowed the saliva to escape from the mouth. Under the repeated application of nitrate of silver and the support afforded by adhesive plaster, this opening was at length permanently closed on November 15th. The reconstructed mouth, as shown by Fig. 24 (taken from a




FIG. 24.

photograph of a plaster cast of the face), presents the upper lip as doubled up on itself, much increased in thickness, protruding, and consequently overhanging the short, tense, retracted under lip. To equalize the dimensions of the two lips, and restore the mouth to its natural angular shape, was the object of the next operation, which was performed on November 30th.

Third Operation.—The method employed was that described on page 22, and was applied first to the left half of the mouth only. A satisfactory result was obtained, and the natural angular shape of the mouth was restored on that side.

Fourth Operation.—On January 4, 1870, the same operation was applied to the right half of the mouth as had been applied to the left. It was feared that sloughing might take place on this side of the face from the presence of numerous cicatricial lines, resulting from the previous operations. None, however, occurred to mar the result of the operation. The apex of the triangular patch, after its transfer from the cheek to the side of the chin, did slough, but without any detriment to the result. By these two operations the mouth was restored to its natural shape and functions, except that there remained a notch on the right half of the under-lip border, near the angle of the mouth, where the saliva escaped uncontrolled. This defect in the result of the last

operation, as shown in Fig. 25 (taken from a plaster cast), was owing partly to a greater deficiency of material on the right side of the chin originally, and partly to the right half of the under lip having united at the symphysis, below the level of the left half, after the first operation. The patient spent the following summer in the country, and returned to




FIG. 25.

the city in October, much improved in health. The parts involved in the previous operations had also improved, having regained a good degree of softness and pliability. The discomfort caused by the escape of saliva demanded an operation for the closure of the notch upon the under lip. It was performed on October 5, 1870, as follows:

Fifth Operation.—The border of the notch was split lengthwise by an incision along its middle, and the edges on both sides of the split were pared off, beveling, so as to increase the thickness of the fresh-cut border. The upper lip was then transfixed a little above the vermilion border, at the middle of

its right half, and an incision carried toward and around the right angle of the mouth in such a manner as to detach a strip of lip border, half an inch wide, that would include the angle of the mouth. This strip, still connected with the upper lip, was brought round lengthwise, and secured to the fresh-cut edge of the notch by fine-thread sutures, inserted close together. Though a material improvement was effected by this operation, the escape of saliva was not entirely controlled, and the mouth was somewhat shortened.

Sixth Operation.—Was performed on April 12, 1871, for the purpose of restoring the angular shape of the mouth on the right side. It was executed according to the method described on page 28. A good result followed, and the natural angular shape of the mouth was restored.

Again the patient spent the summer in the country, and on his return to the city, in September, a further improvement in the condition of his face was observable. The mouth, though restored to its natural angular shape on the right side, was still scanty in length. A shallow notch on the right half of the under-lip border also remained, and allowed some escape of saliva, especially when the head was inclined forward, and his attention was not directed to controlling it. After a little persuasion, seconded by his mother's influence, the patient consented to a

final operation for remedying the defects still existing. It was performed on September 23, 1871, as follows:

Seventh Operation.—The first step of the operation was to lengthen the mouth on the right side by extending the angle further outward. This was accomplished by repeating the sixth operation, as above



FIG. 26.—Condition in March, 1872.

described. The next step was to obliterate the notch on the right half of the under-lip border. To accomplish this the following method was executed: Two incisions, commencing one on either side of the notch, a little below the lip border, and extending through the entire thickness of the lip, were carried

downward in converging lines that met below the chin. The included triangular patch, having for its base the notched portion of the lip border, and retaining its connection for support on both sides of the notch, was pushed upward, and secured on a higher level by bringing together the edges of the wound below the patch, and securing them in contact by a pin suture, wound with cotton yarn. A second pin suture was inserted higher up, and made to traverse both edges of the wound, as well as the patch interposed between them; an additional thread suture below completed the adjustment. Primary union without any suppuration followed, and on the third day the last suture was removed.

As a final result of these several operations, the patient's mouth was restored to its normal shape and dimensions. The saliva no longer escaped uncontrolled. The lips performed their natural motions, and articulation was but little affected. Fig. 26, representing his condition in March, 1872, was engraved from a photograph. At the present time—March, 1873—still further improvement is noticeable. The numerous cicatricial lines, intersecting the surface on both sides of the chin and cheeks, have shrunk to the level of the adjacent surface, and they, as well as the skin itself, are perfectly soft and pliable.

Patient was again seen in January, 1876, and the good results of surgical treatment previously ascer-

tained were found to continue unimpaired, as well in respect of the appearance of the parts themselves as of their functions.

CASE V.—*Reconstruction of the Mouth after Removal of the Under Lip for Disease.*

S. D., aged forty-nine, married, native of United States, a calker by occupation, was admitted into St. Luke's Hospital May 28, 1870, with epithelial cancer of the under lip. The narrative of the case will be restricted to the patient's condition at the time of entering the hospital, which was as follows: A salient, morbid growth involved the middle portions of the under-lip border to within half an inch of both angles of the mouth, and extended downward upon the chin to within a finger's breadth of its lower border. The left half of the tumor stood out prominently, and its everted margin overhung the adjacent sound surface. The upper lip was somewhat folded upon itself and drawn down at the angles of the mouth, in consequence of the removal of a V-shaped portion of the under lip by an operation performed in the month of January preceding, for the removal of a diseased growth involving the lip border, which had preceded the present growth. The wound had been treated after the operation in

the usual manner, but before healing was complete the growth reappeared in the wound, and at length attained the dimensions above described. A glandular enlargement, about the size of a cranberry, existed under the jaw on the right side, at a point below the anterior edge of the masseter muscle. There was also a much smaller one on the left side, in a corresponding situation. (*See Fig. 27, taken from a photograph of a plaster cast of the patient's face.*)




FIG. 27.

First Operation.—Was performed May 29th, as follows: An incision, commencing at a point within less than half an inch of each angle of the mouth, was made to divide the under-lip border, and extend vertically downward on either side of the morbid growth, till they both joined a transverse incision crossing the lower part of the chin, *a* to *b*; *a* to *b*, Fig. 27. The quadrilateral patch thus formed, in-

cluding the morbid growth, was dissected off from the underlying periosteum. The transverse incision was then continued outward on both cheeks to a point within a finger's breadth of the angles of the jaw, and thence extended upward a distance of two inches in a line curving slightly forward (*b* to *c*; *b* to *c*). The cheek flaps thus designed were dissected up on both sides from their underlying connections, and the mucous membrane, lining their inner surface, was alone divided along the anterior margin of the masseter muscle upward, and thence forward along the line of its reflection, from the upper jaw to the cheek, as far as the upper canine tooth. The two cheek flaps, thus freely liberated, were glided forward edgewise toward each other, and made to meet over the symphysis menti, where they were secured in accurate coaptation by three pin sutures and intermediate fine-thread sutures. The spaces left bare on the cheeks by the transfer just described were closed by approximating their opposite edges, and securing them together by sutures, which, owing to the laxity of the skin, was effected without any dissection. The facial artery was unavoidably divided on both sides, and consequently had to be ligated. These, with other ligatures that had to be applied to bleeding vessels, were gathered into a loose skein and brought out at the nearest point of exit. The mouth, as the result of this new adjustment, assumed a cir-

cular and pouting shape, which, however, still permitted liquid nourishment to be taken through a tube and by means of a spoon. In the subsequent progress of the case a moderate degree of inflammatory tumefaction supervened. Saliva escaped for a few days from between the edges of the wound on the right side of the chin. The edges of the cheek flaps, where they met in front over the symphysis menti, and were subjected to a great degree of tension, required the most careful management to secure final union. Fortunately union was secured by primary adhesion at the lip border, but it failed below to the extent of an inch, and here the first pins inserted had to be reënforsed by a succession of fresh pins, and at last by resorting to a beaded-wire clamp suture, inserted in the manner described on page 17; and this first clamp was succeeded by a second one, which was finally removed and dispensed with on June 14th, when union was completed at all points. Fig. 28, which shows the result of this first operation, was taken from a photograph of a plaster cast.

Second Operation.—Was performed on the 23d of June, for the purpose of equalizing the length of both lips, and restoring the natural angular shape of the mouth. It was executed according to the method described on page 22, and was applied to both sides of the face at one operation. Fig. 28 shows with greater exactness the direction of the in-

cisions, as designated by pins inserted erect at selected points and dotted lines drawn between them. For the first few hours after the operation the surface of the transferred patches had a pale, ashy look, which, however, disappeared on the day following without any subsequent sloughing. Primary union took




FIG. 23.

place at all essential points. The reconstructed mouth presented a symmetrical and natural shape, without any interruption of the continuity of the lip border at its new angles. Patient was discharged July 7th, at his own request, and readmitted October 14th, for the removal of a tumor situated in the submaxillary region of both sides; their existence had been recognized before the first operation. They were then of small size, and it was hoped they might disappear after the removal of the diseased under lip.

They had, however, considerably increased in size, and their true character was no longer doubtful. Patient's general health had much improved. His mouth retained its symmetrical and natural shape, both angles being perfect in form. Though somewhat scanty in length, the mouth performed all its functions so satisfactorily that patient was unwilling




FIG. 29.

to accede to the proposal to increase its length by another operation. The tumor on the right side adhered closely to the lower edge of the jaw, and after removal was found of the size of a hickory-nut. That on the left side was loose and movable, and of the size of a cranberry. Both wounds having nearly healed, patient left the hospital, and returned to his home October 21st.

Fig. 29, also taken from a photograph of a plaster cast, shows the final result. The author learned that

the disease reappeared several months subsequent to his leaving the hospital, and progressed to a fatal termination.

CASE VI.—*Reconstruction of the Mouth after the Loss of the Right Half of both Lips; also, Es-march's Operation for Anchylosis of the Jaw.*

GEORGE K., aged six and a half years, of German parentage, resident of Williamsburgh, Long Island, N. Y., of fair complexion and light hair, was admitted into St. Luke's Hospital in May, 1866. The loss of parts sustained in this case appeared, from the father's statement, to have been caused by cancrum oris, that occurred in the progress of typhoid fever; whether it followed the administration of mercury or not could not be satisfactorily determined. His condition, when I first saw him, was as follows: One-half of the upper, and two-fifths of the under lip, with the right angle of the mouth and the adjacent portion of the cheek, had been destroyed, leaving the subjacent teeth and gum-surface exposed. The cicatrized margin of the cheek was retracted, depressed, and closely adherent to the upper and lower jaws, binding them together immovably in close contact. The lining mucous membrane of the right cheek having been destroyed, the cavity of the cheek was oblit-

erated. What remained of the upper-lip border terminated below the septum nasi in a rounded, shrunken, and somewhat everted extremity. The remains of the under lip constituted about three-fifths of its original size, and terminated in a rounded, everted extremity at a point immediately below the right canine tooth of the lower jaw. The columna nasi had been destroyed, leaving the cicatrized inferior border of the septum exposed. In consequence of the closure of the jaw the introduction of food into the mouth could only take place along the cavity of the left cheek, and across the space between the last molar teeth and the ascending ramus of the jaw. Soon after his admission to the hospital a necrosed portion of the lower jaw on the right side was removed. It was found to consist of the entire breadth of the jaw vertically, and included three-quarters of an inch in length of its lower margin, while upon its upper margin there still remained the entire alveolar socket of the second bicuspid tooth, with one-half of the socket of the first bicuspid anterior to it, and one-half the socket of the first molar posterior to it. Notwithstanding the loss of so considerable a portion of the bone, the reproduction of new bone was so complete that no trace of deficiency could be detected by the finger passed over the inferior margin of the jaw. Patient's general health, though pretty good when he entered the hospital, steadily improved

afterward by the aid of generous diet and daily out-of-door exposure. (*See* Fig. 30.)

First Operation.— Was performed June 20th. The right cheek was detached from the jaws above and below, and the dissection continued in every



FIG. 30.

direction, till the jaws could be separated far enough to admit the thumb edgewise between the front teeth. The thinned cicatricial edge of the right cheek, bordering on the region of the angle of the mouth, was pared afresh for adjustment to the new lips. What remained of both lips was now detached from their connections, the upper lip by an

incision commencing at the inferior border of the ala nasi, and dividing the entire thickness of the lip on a line parallel with the lip border, and continued onward to the middle of the left cheek. The lower lip was detached by an incision across the middle of the chin, continued onward, parallel with and as far into the left cheek as the previous incision of the upper lip. The bifurcated quadrilateral flap thus obtained, which consisted of the remains of both lips, and was lined with mucous membrane, was advanced toward the right side of the face, and its two extremities adjusted, with the lip borders in contact with each other, to the edge of the right cheek, already prepared. Pin sutures were employed to secure the ends of the flap in place, and thread sutures, inserted close together, served to secure the upper and lower edges of the flap in their new relations. This adjustment was completed in such a manner as not to have any strain upon the sutures. Warm-water dressings were directed to be applied to the parts, and liquid nourishment to be given through a tube. Moderate inflammatory swelling followed the operation, but it began to subside on the third day. On the fifth day most of the sutures had been removed, primary union having taken place at nearly all points. Patient's appetite was good and his general condition satisfactory.

July 5th.—Healing was complete at all points.

Means were employed after the operation to prevent the closure of the jaws, by keeping wedges of wood between the teeth during the progress of cicatrization. They could, however, be borne only a part of the time, and ultimately proved of no avail. The newly-constructed mouth, as shown by Fig. 31, is



FIG. 31.

scanty in length, and situated mostly to the right of the median line, with its left angle on a line below the left orifice of the nostril. This defect, in the length and symmetry of the mouth, it was proposed to remedy by a second operation, which was performed on the following 23d September.

Second Operation.—Was executed according to the method described on page 28, and promised well. On the fourth day after the operation the last suture was removed. Though a considerable improvement in symmetry was effected by this operation, the mouth was still deficient in length, and scarcely exceeded one inch and a half. In the month of May, 1867, patient, after a long absence at home, was re-admitted into the hospital for a

Third Operation.—This was a repetition of the preceding one, and was performed for the purpose of extending the mouth still further at the left angle. The improvement in the symmetry of the mouth and the general expression of the face, resulting from this operation, was quite satisfactory. Patient still suffered, however, all the discomfort connected with the closed condition of his jaws, and could introduce food into his mouth only between the left cheek and teeth, as already described. The only expedient adapted to remedy this condition was Esmarch's operation for establishing an artificial articulation, at a point anterior to the cicatricial bands which held the jaws in contact. This was resorted to on July 1, 1867.

Fourth Operation.—It will be borne in mind that the cicatricial band holding the jaws in contact occupied the entire region of the right cheek, and terminated anteriorly in a callous edge, extending

between the upper and lower bicuspid teeth. At a point anterior to this callous edge a portion of the lower jaw was to be excised, for the purpose of establishing an artificial articulation. All the muscles concerned in depressing the lower jaw, having their insertion on either side of the symphysis, would be undisturbed, and consequently would continue to perform their office in acting upon the jaw after the proposed operation. The procedure was as follows: An incision was made along the inferior edge of the jaw down to the bone, from a point near the angle to a point three-quarters of an inch distant from the symphysis. The outer and inner surfaces of the jaw were denuded to the same extent. The bone was then perforated by a drill of the size of a goose-quill on a line below the first bicuspid tooth, to facilitate the division afterward of the entire bone by the action of a strong, cutting bone-pliers. The same procedure was applied posteriorly on a line below the second molar tooth, and the included fragment of bone, measuring more than one inch in length, was removed. Special care was taken to avoid the facial artery by drawing it out of the way posteriorly. A mass of callous tissue, in which the upper teeth were imbedded, was pared away. The cut ends of the bone were gnawed smooth with Lürer's rongeur forceps. The anterior liberated portion of the lower jaw could now be separated from the upper jaw so

as to admit a finger edgewise between the molar teeth on the left side. A tent of lint, of the size of the little finger, was inserted, with one end passing out at the right angle of the mouth, and the other through the wound below the jaw. The remainder



FIG. 32.

of the wound was closed by sutures. Water dressings were applied to the face and neck. The subsequent inflammatory action was moderate, and soon began to subside. The final result of this operation was as follows: The space left after the removal of the portion of jaw-bone diminished by the gradual approximation of the opposite ends of the bone.

The degree of mobility, however, of the anterior fragment of the jaw was only limited, but yet it sufficed to permit the introduction of food into the mouth directly between the front teeth. (*See Fig. 32.*)

The author regrets that he has not been able to discover the whereabouts of this patient, so as to ascertain his condition after the test of a long interval of time.

CASE VII.—*Mutilation and Distortion of the Mouth from a Shell-Wound.*

EGBERT H., aged twenty-two, native of Vermont, a private soldier in Company C, Sixth Vermont Regiment, Sixth Army Corps, received a wound on September 19, 1864, at Winchester, Va., from a fragment of a shell, that laid open the right cheek from the angle of the jaw to the mouth, lacerated the under lip at the right angle of the mouth, and carried away the upper and lower front teeth. The nose and upper lip were also split vertically, and the under jaw was fractured at the symphysis, with comminution of the bone. The parts having been healed for some time, the condition of the face, at the time of the operation, was as follows: The mouth was shrunk and contracted to half its natural size, and both lips shortened. The right half of the under-lip border

overlapped and adhered to the alveolar border of the jaw, from which the teeth were absent. A deep notch in the lip, capable of lodging the little finger, had resulted from this adhesion, and permitted a constant escape of saliva from the mouth, to the great annoyance and discomfort of the patient. The surface of the skin bordering on the upper lip and right angle of the mouth was wrinkled and cicatricial. A conspicuous scar crossed the right cheek from the angle of the jaw to the mouth. The adhesion of the right half of the under lip to the jaw, though it involved the entire depth of the lip, did not spread out laterally. The loss of bone, consequent on the comminuted fracture of the lower jaw at its symphysis, had been followed by an approximation of the two halves of the jaw to each other, thereby narrowing the dental arch so that the upper and lower teeth of both sides could not be brought simultaneously into contact with each other. Owing to this deformity mastication and articulation were somewhat affected. Patient wore habitually a small tin gutter, to catch the constant flow of saliva. (*See Fig. 33.*)

First Operation.—Was performed October 28, 1864, at the Central Park Hospital, at the request of Surgeon Clements, U. S. A., in charge, and in the presence and with the aid of the medical officers of the hospital, as follows: Two incisions, dividing the

under-lip border, were commenced, one at the right angle of the mouth, *a*, the other at the middle of the lip, *b*, and continued downward, in gradually-converging lines, to a point, *c*, under the chin. The V-shaped patch thus formed, including as it did the notch upon the lip border and the adherent portion




FIG. 33.

of the lip below it, was dissected off from the underlying periosteum. The remaining left half of the lip and the adjacent cheek were detached from the jaw as low down as its inferior border, and as far back as the last molar tooth, after the buccal mucous membrane had been first divided to the same extent

along the line of its reflection from the jaw to the cheek. This procedure permitted the detached parts to be glided toward the right side of the chin, and thus to contribute to close the space left by the removal of the V-shaped patch of lip. In order to obtain additional material for the same purpose on the right side of the chin, a transverse incision, *d* to *a*, was carried through the right angle of the mouth across the cheek to within a finger's breadth of the anterior border of the masseter muscle, and thence a second incision, *a* to *e*, was extended downward and a little forward to a point below the edge of the jaw, on the same level with the apex of the V-shaped bare space under the chin. The quadrilateral patch thus formed, having its upper half lined with mucous membrane, was dissected up from the jaw, but left attached below it. It was then glided forward edgewise till it met the left half of the under lip, and their confronted edges could be adjusted to each other by pin sutures, and fine-thread sutures in the intervals between the pin sutures. By this adjustment the deficiency of the under lip was supplied, and the lining mucous membrane of the quadrilateral patch made to confront the surface of the jaw, denuded of its mucous membrane by the previous removal of the adherent portion of the lip. New lip border was constructed on the upper edge of the transferred cheek patch by excising a prism-shaped

strip of tissue from between the skin and mucous membrane, and, after lapping the latter over the former, securing them accurately together by fine sutures, inserted close to each other. As it was necessary to lengthen the mouth on the right side beyond its previous limits, an addition of three-fourths of an inch had to be made to the upper-lip border, which was done in the manner just described in its application to the under lip. A new angle was also constructed for the mouth by securing the opposite edges of the divided cheek together at a point where the newly-constructed upper and lower lip borders terminated. The reconstruction of the mouth being completed, there still remained a triangular-shaped space of raw surface left bare upon the right cheek by the transfer of the quadrilateral patch. To provide a covering for this space the transverse incision, which had divided the cheek to within a finger's breadth of the edge of the masseter muscle, was continued in the same line through the skin alone to the distance of one inch, and the skin bounding the raw surface posteriorly was then dissected up and stretched forward so as to meet the edge of the advanced cheek patch, and thus cover the bare surface. This adjustment was secured by the requisite number of thread sutures. The operation thus completed occupied three hours, including the interruptions necessary to maintain the anæsthesia. The inflamma-

tory tumefaction succeeding the operation was considerable for the first three days, and then began to abate; the febrile reaction was moderate. Suppuration took place only at the lower angle of the wound under the chin, and was of short duration. The management of the sutures was the same as has already been sufficiently noticed. Toward the end of the second week patient left his bed and went about the ward. As a result of the operation patient's appearance was much improved; the saliva no longer escaped from the mouth uncontrolled, and both mastication and articulation were benefited. Certain defects, however, still remained: the mouth was one-sided, three-fifths of its length being situated to the right of the median line, and two-fifths to the left. The lips being without support, in consequence of the absence of front teeth, retracted toward the cavity of the mouth. Patient being anxious to have these defects remedied, if possible, engaged to return to the city for another operation, after a visit to his family. On his return to the city he was admitted into the New York Hospital, December 12, 1865.

Preliminary to a second operation, Mr. J. A. Bishop, a skillful dentist, of No. 34 East Twenty-first Street, kindly undertook to adapt to the patient's mouth a plate, made of vulcanite, with artificial teeth in front, to supply the deficient teeth and afford support to

the lips. Mr. Bishop's success was complete, and the result satisfactory.

A Second Operation.—Was performed January 9, 1866, for the purpose of lengthening the mouth at the left angle, and thereby restoring its symmetrical shape. It was executed according to the method described on page 28. A good result followed, alike satisfactory to the surgeon and patient. Fig. 34,



FIG. 34.

which shows the final result, was engraved from a photograph taken before patient's discharge from the hospital, January 19, 1866.

CASE VIII.—*Loss of a Large Portion of the Lower Jaw-bone, with Extensive Mutilation of the Face and Distortion of the Mouth, produced by a Shell-Wound.*

WM. SIMMONS, aged twenty-one, a private soldier of Company 1, New York Heavy Artillery, was admitted into New York Hospital, October 26, 1865. On the 25th of March preceding, while serving with the army before Petersburg, Va., patient was struck on the right side of the face by a fragment of shell, that lacerated the cheek extensively, and carried away the body of the lower jaw. The condition of the face, after the injured parts had been healed for two months, was as follows: The body of the inferior maxilla having been carried away, the chin, for lack of support, had lost its prominence and was sunk. The entire right ramus of the jaw, including the angle and alveoli supporting the last two molar teeth, remained *in situ*, while on the left side the upper half of the ramus only was preserved. From the middle of the zygoma on the right side a linear cicatrice extended obliquely downward and forward upon the cheek to a deep depression at the angle of the mouth, where the extremity of the upper lip and neighboring parts adhered closely to the underlying alveolar border of the upper maxilla, from which the teeth had been carried away. The upper-lip border

retained its normal horizontal position, but its right half was increased in length, in consequence of the adhesions at its extremity. The right half of the under lip, having been completely detached from its connections by a lacerated wound crossing the middle of the chin transversely, had, in contracting new adhesions, settled below its proper level, leaving a space of one finger's breadth between the right halves of the two lips, from which saliva escaped constantly uncontrolled, to the great discomfort of the patient. The surface of the left side of the chin was uneven and intersected by cicatricial lines. On the inside of the mouth, behind the under lip, a callous ridge stretched obliquely across the floor of the mouth, and terminated on the left side at, and adhered to, the extremity of what remained of the ramus of the jaw. This band, which formed a part of the cicatrice uniting the under lip to the chin, appeared to serve as a substitute for the lost bone in front, and afforded a firm support for the anterior attachments of the tongue. The last upper molar tooth alone remained *in situ* in the upper jaw on the right side, while the canine and all its fellows posterior to it remained *in situ* on the left side. By the aid of the finger, introduced into the mouth, the under surface of the right half of the tongue was found adherent to the opposite surface of the floor of the mouth, and the protrusion of the tongue was thereby rendered impossi-

ble. His food was restricted to soft solid and liquid articles of nourishment. Deglutition was unimpaired, but articulation was so defective that he was averse to holding conversation, and preferred to communi-



FIG. 35.

cate with others by signs and the use of a pencil and paper. His appearance was ruddy, and his general health good. (Fig. 35.)

Operation.—Performed November 7th, as follows: The under lip was detached by a transverse incision carried across the chin above the cicatricial line connecting the lip with the chin, to a point one finger's breadth below the left angle of the mouth. The lip

border, which before being detached had assumed a fan-shape, could now be straightened out and applied to the upper lip throughout its entire length. In order to reconstruct the right angle of the mouth, the upper lip was detached at its right extremity from its adhesions to the upper jaw. A point was then chosen on its border, at such a distance from the median line as would make the two halves of the lip equal in length, and at this point the border was pared away obliquely. At a corresponding opposite point upon the under lip the border was also prepared in the same manner, and the two lips were then confronted and secured together at their fresh-cut edges by a single pin suture and additional thread sutures. The next step of the operation was to cover up the depressed cicatricial line upon the cheek which was so conspicuous. It was accomplished by carrying two parallel incisions, one on either side of the cicatricial line, downward and forward to the angle of the mouth. The edges of these two incisions were then dissected up, and brought together and secured by sutures, so as to cover up the cicatrix, after its surface had first been pared and made raw. This adjustment was extended to the newly-reconstructed right angle of the mouth. The same procedure was then applied to the chin. An incision was carried across the chin below and parallel with the cicatricial line already mentioned, to a point below the

left angle of the mouth. The lower edge of this transverse incision was next to be adjusted to the inferior cut edge of the detached under lip, so as to cover up the cicatrix intervening between them. In order to effect this adjustment the lower edge of the transverse incision required to be everted, which could only be effected after carrying a vertical incision from the terminus of the transverse incision below the left angle of the mouth, a distance of one inch and a half upon the neck. In the course of this vertical incision a flattened cyst, of the size of a silver half-dollar piece, filled with brownish viscid fluid, was encountered under the skin, and removed entire. The angle of skin included between the last two incisions was dissected up, and its transverse edge made to overlap and cover the cicatricial line, and meet the inferior border of the detached under lip; the two were adjusted to each other by sutures. Pin sutures were employed in this adjustment at selected points where the greatest support was required, and between them thread sutures were added. The reconstructed mouth had now assumed its normal shape and dimensions with both lips in contact. No adhesive plaster was applied. Warm-water dressings were directed to be kept upon the part, and liquid nourishment allowed to be given through a tube.

8th. — Progress favorable. Inflammatory tumefaction not excessive; febrile reaction moderate.

Changed the yarn on the pins, and removed some of the alternate thread sutures.

9th.—Doing well. Changed yarn; removed additional sutures, where they could be dispensed with.

10th.—Primary union secured at nearly all points. Removed all the pins and most of the remaining



FIG. 36.

thread sutures. A free suppuration escaped from the lower angle of the wound, below the chin. A superficial slough formed over the right zygoma, but it in no way marred the result of the operation. Strips of adhesive plaster were now applied at different points to support the newly-healed parts. The suppuration gradually diminished, and at length

ceased, and healing was finally completed at all points. The lips, restored to their natural relations to each other, performed their functions, and controlled the escape of saliva. Some improvement in articulation and a marked improvement of the general expression of his countenance were observable. On the 12th of December following patient returned to his home in the country. In May, 1866; he revisited the hospital. A further improvement had taken place in his general appearance, and especially in his articulation, which had now become so much more distinct that he no longer shrunk from engaging in conversation, having discarded entirely the use of pencil and paper in communicating with others. The result is shown by Fig. 36.

CASE IX.—*Closure of an Opening into the Superior Meatus of the Right Nasal Fossa.*

MARGARET K., aged twenty-one, native of Ireland, unmarried, was admitted March 3, 1869, into the New York Hospital, and gave the following statement of her case: After passing through an attack of scarlet fever when nine years old, a black spot appeared on the right side of the upper part of her nose, and was the occasion of the opening which still exists. Its situation and shape are accurately repre-

sented in the accompanying Fig. 37, taken from a photograph. It is of an ovoid form, more than one inch in its vertical diameter, and three-quarters of an inch in its greatest transverse diameter. It communicates directly with the right nasal fossa, and allows a free passage to the air. Its outer edge approaches close to the inner canthus of the right eye.



FIG. 37.

The skin surrounding it is thin, pliable, and somewhat overlaps its margin; the inner surface of the cavity, as far as it is visible, appears healthy. Her voice is unaffected. A patch of adhesive plaster is habitually kept applied to the opening to conceal it. Patient being naturally anxious to be rid of so con-

spicuous a disfigurement, consented willingly to an operation for its removal.

Operation.—Performed March 5th, as follows: The skin at the margin of the opening was dissected up and everted, a procedure requiring great care, owing to the thinness of the skin, especially where it bordered on the inner canthus of the eye. A pattern of the size of the opening, cut from oiled-silk, served as a guide in shaping the patch of skin which was to be raised from the forehead. To prevent this patch from becoming too much twisted, when transferred to its new locality, its position on the forehead was so chosen that its long axis inclined obliquely at an angle of forty-five degrees toward the opening, and its pedicle of attachment lay above and close to the inner extremity of the left eyebrow. After dissecting up the patch of skin designated by the letters *b, a, c*, a strip of skin, *e, d*, intervening between it and the opening, was also dissected up, to make room for the transfer of the patch, but left attached above the right eyebrow, and reserved for subsequent use. After adjusting the patch to the opening by sutures, the reserved strip of skin was utilized to cover the surface left bare upon the forehead, which it did so completely that no bare spot was left uncovered after the completion of the operation. Warm-water dressings were directed to be applied to the parts.

The subsequent progress of the case was favorable, and primary union followed at all points, except at that portion of the circumference of the patch which bordered on the inner canthus of the eye; here union failed to take place, in consequence of the opposite edges becoming inverted. To remedy this defect a second operation was performed April 6th.

Second Operation.—The ununited edges were pared afresh, and two pin sutures inserted, special care being taken to evert the edges and maintain



FIG. 38.

their cut surfaces in contact. Additional thread sutures were also necessary to complete the adjustment. Perfect union followed this operation. The trans-

planted patch, from some excess in size, formed in its new locality a rather conspicuous bulging ridge, particularly between the eyebrows, where its pedicle had undergone a twist. To remove this disfigurement a third operation was performed April 17th.

Third Operation.—A prism-shaped strip of skin was excised from the ridge along its entire length, and the opposite edges of the wound brought together into exact coaptation, and secured with numerous fine-thread sutures. Primary union followed, and a level surface remained. The final result was a decided improvement in the appearance of the face. (*See Fig. 38.*)

CASE X.—*A Rhinoplastic Operation for the Restoration of the Apex Nasi after it had been bitten off.*

W. W. G., aged thirty-five, a resident of the city, was brutally assaulted in the evening of May 12, 1872, and during the affray had the apex of his nose bitten off by his assailant. The condition of the parts, twelve days after the occurrence, was as follows: All inflammatory swelling of the nose had subsided. A healthy suppurating surface indicated the extent of the lost parts, which included the skin covering the apex, and adjacent ridge of the nose as high up as its middle, and also both sides of the nose

to a point within half an inch of the junction of the nose with the cheeks. The entire denuded surface was equivalent to about one-third of the superficies of the organ. The alæ nasi were disconnected anteriorly, and both had sustained about an equal amount




FIG. 39.

of loss of substance. The columna remained entire. The ridge of the cartilaginous septum was denuded, but had sustained no loss of substance. (*See* Fig. 39.)

Operation.—On Saturday, May 25th, an operation was performed at patient's residence, with the aid of Prof. A. C. Post, M. D., and Drs. T. E. Satterthwaite, J. N. Beekman, and N. S. Westcott, as fol-

lows: The anterior edges of what remained of both alæ were pared and made straight, and an incision was then carried upward on both sides of the nose, on a line continuous with these edges, as high as the inner extremities of both eyebrows. The patch of skin included between these two incisions was dissected up from the dorsum nasi and between the eyebrows, and left connected above. A pattern, of the shape of the entire denuded surface, cut from oiled-silk, was laid upon the forehead in an inverted vertical position, immediately above the inner half of the right eyebrow, with one edge bordering on the incision that terminated at the inner end of the eyebrow. The patch of skin underlying the pattern having been outlined by an incision, was then dissected up from the perieranium, but left connected below at the margin of the orbit. Suitable allowance had to be made for shrinkage of the patch, especially in the direction of its length, after its transfer.

The skin, previously displaced from the nose and from between the eyebrows, was now dissected up still further toward the left side of the forehead, in order to afford a sufficient extent of raw surface, with which the under surface of the forehead patch might come into immediate contact after being transferred to its new locality. The patch of skin from the forehead was now brought around edgewise from right to left, and from above downward, till it reached its

destination, and was spread out upon the nose, and made to extend downward beyond the apex and overhang it. Special care was taken that there should be no strain at the pedicle of the patch, where it was doubled upon itself, and where any obstruction of the circulation would endanger its vitality. In order to confront more exactly the edges of the patch with the opposite edges of the space it was to occupy, the latter were dissected up and everted sufficiently for the purpose. Pin sutures were inserted at select points, and fine-thread sutures in the spaces between them, to secure the adjustment. The patch of skin displaced from the nose and between the eyebrows was now carried upward, and employed to cover the surface on the forehead left bare by the transfer of the forehead patch. It proved sufficient to fill up the lower half of the space, and was secured there by sutures. The remaining upper half of the bare surface, which encroached upon the hairy scalp, was covered with a collodion crust (*see* page 13). Both patches of skin, in being transferred, were necessarily doubled upon themselves, and formed two prominent folds of a flattened conical shape, standing out on the surface, one being situated above each eyebrow. (Fig. 40, *a* to *b*.) But few ligatures were required to secure bleeding vessels, and these were brought out at the nearest point of exit. Wet applications were avoided, and the parts covered with

a double thickness of sheet-lint, spread on one side with cerate to prevent its adhering. The operation occupied two hours, and was well borne by the patient. His subsequent progress was favorable and requires no special notice. On the fourth day the last sutures were removed. On the sixth day the




FIG. 40.

collodion crust became detached from the forehead and came off, and the underlying healthy, granulating surface was thereafter dressed in the ordinary way, till it finally cicatrized. The result of the operation is shown by Fig. 40.

Second Operation.—Was performed June 28th, for the purpose of leveling the two prominences re-

maintaining upon the forehead, one being situated over each eyebrow. The procedure was as follows: A curved incision was carried half around the base of each prominence on its broadest side, and the prominence itself raised up from its underlying connections; and split across its middle. After being unfolded and spread out, the redundant skin was pared



FIG. 41.

away, so as to be level with the surrounding surface, after the edges had been adjusted to each other and secured by sutures. This operation did well, and was followed by the desired result. The end of the patch of skin covering the nose, which overhung the

apex, was trimmed and shaped after having shrunk as much as it would. Fig. 41, showing the final result, is from a photograph, taken on the succeeding 12th of October, when the following particulars were noticed: Sensation, which for several weeks following the first operation, continued to be referred to the forehead when the surface of the patch on the nose was irritated, had regained its normal condition, and was referred to the actual seat of irritation. The principal cicatrice upon the forehead, where the patch of skin had been removed, had shrunk to a small size, and was concealed by the hair. The other cicatrices were only linear and not conspicuous. The lower third of the dorsal surface of the nose having been derived from the scalp, continued to yield its hairy growth, and required to be shaved at short intervals. Patient was seen as late as October 14, 1875, when all the parts involved in the operation above reported continued in good condition.

SECOND CLASS.

CONGENITAL DEFECTS FROM ARREST OR EXCESS OF DEVELOPMENT.

By far the most numerous of this class are cases of harelip. Their successful treatment depends so much on the proper execution of the several details which make up the operation, such as the management of the patient, the preparation of the parts for readjustment, the choice of sutures, the manner of inserting them and their subsequent management till healing is completed, that, in order to avoid repetition, some general remarks on these several topics will first be presented.

1. *The Management of the Patient.*—In the case of a child the arms should be brought down to the sides of the body, and secured by a folded napkin passed around them and fastened with pins. The head should be held steadily in position between the hands of an assistant. In the administration of anæsthetic, after its full influence has been produced, its continued action during the progress of the operation may be most conveniently maintained by the

use of a small sponge, held in a dressing forceps, and applied to the nostrils of the patient.

2. *Preparation of the Parts for Readjustment.*— In order to hold the parts on the stretch, and thereby enable the operator to make the requisite incisions with the utmost precision, the following expedient will be found very satisfactory: A needle, armed with a coarse thread, is to be passed through each half of the lip at the angle bounding the cleft on either side, a point being chosen for the passage of the needle that will not interfere with the subsequent incisions. The ends of each of the threads when tied together will form loops with which to hold the parts on the stretch when required. Each half of the lip is then in turn to be held on the stretch and drawn away from the jaw, while the lip and cheek are detached from the upper jaw by an incision of the buccal mucous membrane, carried along the line of its reflection from the jaw to the lip and cheek as far back as the molar teeth, if necessary. The dissection is also to be extended upward toward the orbit on the level of the periosteum. This procedure, after being applied on both sides, will allow the two halves of the lip to be approximated, and the confronted edges of the cleft to be secured in contact, without there being any strain upon the sutures that are to hold them together. The opposite edges of the cleft are next to be pre-

pared as follows: Each half of the lip, being again held on the stretch, is to be transfixed near its angle by a Beer's cornea knife, and an incision carried upward, along the border of the cleft, as high as, and if necessary into, the nostril. The strips thus detached from both borders of the cleft, but left attached at the angles below, are to be brought down with their fresh-cut surfaces facing each other, and both transfixed with a threaded needle. The ends of the thread tied together afford a loop with which the strips are held evenly on the stretch, and the fresh-pared edges of the cleft at the same time made to confront each other, while one pin suture is inserted close to the vermilion border below, and another close to the columna nasi above. Two or three thread sutures are to be inserted between the pin sutures. The two detached strips of cleft border, still connected at the angles, are now to be severed by dividing them, not at right angles across their length, but obliquely, so that, after their cut surfaces are confronted and united by three fine-thread sutures, they will form a projection standing out beyond the line of the lip border. By this latter expedient (which is credited to Malgaigne) the formation of a notch at the point on the lip border where the two halves unite is best prevented. The advantage of employing Beer's cornea knife, as recommended above, is, that with it the lip can be more

9

readily transfixed and the sections completed with greater precision. When harelip is complicated with




FIG. 42.

a cleft of the dental arch, the cleft divides the arch into two segments of unequal length, and the anterior

extremity of the longer segment projects in advance of the natural curve of the arch, forming a prominence which interferes with the approximation of the two halves of the lip. This prominence requires to be broken down and reduced into position, an operation best effected by means of Butcher's (of Dublin) bone-pliers, an instrument devised for the purpose (*see* Fig. 42). In employing it the bony prominence is to be seized somewhat crosswise between the two blades, with the bent blade applied upon the anterior surface, high up toward the nostril. On pressing the blades together, the end of the bent blade sinks into the bone, and, while the instrument is acted upon by means of the handles from before backward, a fracture is produced and the prominence reduced into its place, where it fills up the notch, and may be made to adhere permanently, if the opposite edges have been previously pared and made raw for the purpose. Consolidation follows without exfoliation of bone or suppuration.

3. *Sutures and their Management.*—This subject has already been fully treated in an introductory chapter, to which the reader is referred.

HARELIP.

The following classification embraces all the varieties of this deformity, of which examples are to be found in the subsequent pages.

First Class.—Single cleft of the upper lip alone, involving most frequently its left half, immediately below the nasal orifice.

Second Class.—Double cleft of the lip alone, involving sometimes only its border; more frequently, however, its entire vertical dimensions.

Third Class.—Single cleft of the lip, complicated sometimes with cleft of the dental arch alone; more frequently with cleft of the bony and soft palates, in addition to the dental arch.

Fourth Class.—Double cleft of the lip, complicated with cleft of the bony and soft palates, and the coexistence of an intermaxillary bone.

To avoid repetition, it may be here stated that all the cases of harelip about to be narrated were operated on according to the method described above. A rare instance of the hereditary development of harelip will be first reported.

A Harelip Family.—Mrs. Molinieri and her three children, all girls, natives of Genoa, Italy, were admitted into St. Luke's Hospital, January 10, 1871. Mrs. M. herself bore the marks of a successful opera-

tion for harelip performed in childhood. She had a brother and sister with harelip; and, besides her three living children, she had had four others, who had all died in early infancy. Three of them had harelip, and the fourth one only was a perfect child. In other words, there were nine instances of the deformity in two generations of a single family.

CASE XI.—*Single Harelip.*

MARIE ANNE, the second child of this family, aged four, had a single cleft, involving the left half of the lip. The left border of the cleft was vertical



FIG. 43.

in its direction; the right border slanting, and diverging from the left (*see* Fig. 43). Operated on January 21st. Primary union followed, and the last

suture was removed on the fourth day after the operation. Strips of adhesive plaster were continued a few days longer, to support the new ad-



FIG. 44.

hesions and relieve them of all strain. Fig. 44 shows the result, without any notch remaining at the lip border.

CASE XII.—*Double Harelip.*

ROSE, the infant, twelve months old, with double cleft of the lip, separated by a central tongue-shaped portion. The clefts involved only about three-fifths of the depth of the lip, and did not quite extend to the orifices of the nostrils above (*see* Fig. 45).

Operated on January 21st. The borders of the central and lateral portions of the lip, after having been pared, were adjusted together and secured by a pin, traversing all the three portions at their upper part, and by fine-thread sutures in addition. Some difficulty was encountered in reducing the central piece



FIG. 45.

to the same level with the lateral pieces, and was not entirely overcome. Primary union failed to take place, except at the vermilion border. By the employment of adhesive plaster, renewed daily, with great care, union by granulation was at length obtained at all points, and completed on the twelfth day. The elevation of the central piece above the

level of the lateral pieces persisted to a slight degree, but might be expected with time gradually to



FIG. 46.

diminish. Fig. 46 shows the result, with only a slight notch remaining at the lip border.

CASE XIII.—*Single Harelip, with Cleft of the Dental Arch and Bony and Soft Palates.*

JACINTA, the eldest child, aged seven. In this case the cleft of the lip involved its left half; the right and largest segment of the alveolar arch terminated anteriorly, at the margin of the cleft, in a rounded prominence, which stood forward in advance of the curve of the arch, and, being uncovered by the

lip, formed a conspicuous feature in the disfigurement of the face. The left ala nasi, being unsupported, was retracted toward the cheek, so as to flatten the nose on its left side (*see* Fig. 47). Operated on Jan-



FIG. 47.

uary 31st. The bony prominence formed by the anterior extremity of the right segment of the alveolar arch was first broken down and reduced into position by the application of Butcher's bone-pliers, as described on page 129. The prominence, after it was reduced, bridged over and filled up the cleft in the alveolar arch. By previously paring the confronting edges bony consolidation was secured. The removal of this prominence also facilitated the approximation of the two halves of the lip and their adjustment to each other. Primary union was secured only at the

lip border. It therefore became necessary to hold the two halves of the lip in contact while union by granulation was taking place. This was effected by the employment of a beaded silver-wire suture, which was inserted high up toward the nose, and at a distance of nearly one inch on either side of the confronted edges of the cleft. The clamp suture was left *in situ* five days, and healing was complete at all



FIG. 48.

points at the end of three weeks. A tooth, growing out of the bony projection that had been reduced, became loose and was extracted. The fractured parts became consolidated without exfoliation or suppuration. A plug of sponge was worn in the left nostril, to keep it distended and in good shape. No notch remained at the lip border. Fig. 48 shows the result.

Other examples of single harelip uncomplicated:

CASE XIV.—*Single Harelip on the Right Side.*

JOHN BLACK, aged twelve, from Piermont, on the Hudson, entered St. Luke's Hospital, January 18, 1871. The cleft divided the right half of the lip, and extended upward, as a shallow furrow, along the floor of the right nostril. The two halves of the



FIG. 49.

lip were but little separated from each other. The right middle incisor tooth occupied the cleft, and stood out conspicuously in advance of its fellows, owing probably to the lack of support from the lip (Fig. 49). Operation performed January 24th, and followed by primary union. The last sutures were

removed on the fourth day. Adhesive straps were renewed daily for a week longer, when healing was



FIG. 50.

complete. A very slight notch remained at the lip border. Fig. 50 shows the result.

CASE XV.—*Single Harelip.*

A FEMALE infant, five months old, from Hudson, Columbia County, N.Y. The cleft, as usual, involved the left half of the lip (*see* Fig. 51). The operation was performed October 12, 1870, and followed by primary union, without any remaining notch at the

lip border. She returned to her home October 30th. Fig. 52 shows the result.



FIG. 51.



FIG. 52.

CASE XVI.—*Single Harelip.*

A MALE infant, three months old, from Hudson City, N. J., with single cleft of left half of the lip (see Fig. 53). Operated on June 11, 1873. Primary



FIG. 53.



FIG. 54.

union followed. Resumed nursing on the seventh day; returned home on the tenth. No notch remaining at the lip border. Fig. 54 is from a photograph taken in August following.

CASE XVII.—*Double Harelip, complicated with Cleft of the Bony and Soft Palates, together with the Presence of an Intermaxillary Bone.*

HARRIET Q., aged eight, from Winsted, Conn., entered New York Hospital, March 16, 1865, with the above deformity of congenital origin. A wide cleft divided, in an antero-posterior direction, the dental arch and bony and soft palates into equal halves. The bony septum narium traversed the entire length of the cleft in the median line. Its inferior border was thin and rounded posteriorly, but expanded out anteriorly into an irregular-shaped mass, which constituted the middle portion of the alveolar arch, and supported the two middle and left lateral incisor teeth, standing out irregularly from its surface. This projecting bony mass, known as an intermaxillary bone, was somewhat deflected from the median plane to the right, so that its right margin touched the anterior extremity of the right segment of the dental arch, thereby obstructing the view into the cavity of the nostril of that side, while it exposed more

fully to view the left nasal cavity. The upper lip was cleft below both nasal orifices, and the two halves of the lip on either side, with their attached alæ nasi, yawned wide apart, thereby increasing the breadth of the nose, and giving it a very flattened appearance. A central tongue-shaped portion of the upper



FIG. 55.

lip was interposed between the two clefts, and rested upon the surface of the projecting intermaxillary bone. It was continuous with, and, as it were, suspended from the columna, which was scarcely half an inch in length. The characteristic defect in articulation which usually attends this condition existed in a marked degree. (*See Fig. 55.*)

Operation.—Performed March 18th, as follows: The projecting intermaxillary bone was excised with bone-forceps, on a line with the inferior border of the septum, after first dissecting up from its upper surface the tongue-shaped portion of lip, as high as its junction with the columna, to which it was left attached. This latter was then brought down into contact with the fresh-cut edge of the septum, and adapted to it by squaring its edges and securing them by sutures to the mucous membrane on both sides of the septum. By this means the scanty columna was lengthened out to its full dimensions. The two halves of the lip were then detached extensively upward and outward, on both sides, from the anterior surface of the upper maxilla, to facilitate their approximation to each other in the median line. In order still further to facilitate their approximation, and also to narrow the cleft in the dental arch, and at the same time restore its natural curve, the projecting ends of its two lateral segments, bordering the cleft on either side, were broken down and reduced into position by the application of Butcher's bone-pliers, as described on page 129. The opposite edges of the cleft in the lip were then prepared and adjusted to each other by sutures, in the manner described on page 126. After the completion of the operation, warm-water dressings were directed to be applied to the parts, and appropriate liquid nourish-

ment, with stimulants in moderate quantity, to be given. The subsequent progress and management require no detailed description. On March 5th the last sutures were removed, and adhesive-plaster applied to support the newly-united parts.

April 29th.—Patient returned to her home in the country. The only remaining incisor tooth in the



FIG. 56.

upper jaw had to be extracted, owing to ulceration of the opposite surface of the upper lip from pressure against it. As a final result of the operation, the dental arch in front was restored to its natural curve, and bony consolidation took place without exfoliation or suppuration. The cleft involving the alveolar arch anteriorly had become narrower. A

slight notch remained at the lip border, where the two halves had united. Fig. 56 is from a photograph taken more than one year after the operation.

THE following are examples of cases of harelip, operated on a second time for the purpose of remedying the imperfect results of previous operations performed in infancy.

CASE XVIII.—A daughter of Mrs. S., aged six, from Cold Water, Mich., had been operated on, in



FIG. 57.

infancy, for single cleft of the right half of the upper

lip. As the result was very imperfect, the parents were anxious to have the defect remedied, if possible, by a second operation. (See Fig. 57.)

Operation.—Performed October 12, 1871. A good result followed after primary union, and without any remaining notch at the lip border. Healing was complete on the sixth day. The result was especial-



FIG. 58.

ly gratifying, from the fact that other surgeons, whom the parents had consulted, had counseled against an operation. One year later the mother wrote: "Hardly any trace now remains of the scar, and we have no doubt she will entirely outgrow it." Fig. 58 shows the result.

CASE XIX.—A physician's daughter, aged sixteen, from Savannah, Mo., had been operated on, when five months old, for single harelip, involving the right half of the lip, with a very unsatisfactory result, as shown by Fig. 59. In order to remedy the



FIG. 59.

defect, a second operation was performed May 27, 1873, and followed by primary union and complete healing on the sixth day. On the tenth day she left for home with her father.

Under date of August 23d following, her father, in a letter inclosing her photograph, from

which Fig. 60 was taken, writes, "The condition of her mouth is still improving."



FIG. 60.

CASE XX.—Miss C., of New York, aged thirty, was operated on, when seven months old, for single harelip, complicated with cleft of the bony and soft palates, involving the left nasal cavity, but without extending through the dental arch anteriorly. The restoration of the upper lip alone was attempted, but the result of the operation was unsatisfactory. A conspicuous notch remained at the border of the lip, and the union between the two halves of the lip was incomplete at the upper part, so that the outer

orifice of the left nostril extended considerably below the level of that of the right. The dental arch, not being involved in the cleft, retained its natural curve. (See Fig. 61.)



FIG. 61.

Operation, May 14, 1874.—After separating the two halves of the lip with scissors, applied exactly on the line of the cicatrice uniting them, the fresh-cut edges were prepared for readjustment to each other by the same method as would be employed in a first operation for harelip (see page 126). Special care was requisite to secure adhesion of the opposite edges of the cleft at its highest point, where it in-

volved the orifice and floor of the nostril, and where union had failed to take place after the first operation. After the opposite edges of the cleft within the orifice of the nostril had been pared and made raw, a beaded silver-wire suture was employed for the purpose of maintaining them in close contact after they were confronted. The wire was made to traverse the confronted raw edges, and was armed at one end with a bead, resting against the septum nasi, low down in the right nostril, and at the other end with another bead, resting against the left cheek, where it joins the ala nasi. The adjustment of the two halves of the lip below and at its border was completed in the manner already described (*see* page 127). The details of the subsequent progress and treatment it is unnecessary to relate, except that the beaded suture was left *in situ* till the eighth day, and performed a useful service in securing adhesion at the point where it was most difficult to effect it, and where we most feared that it would fail. Prior to the operation just described there existed a want of symmetry in the mouth. The portion situated on the right side of the median plane exceeded in length that on the left side. The difference was somewhat increased, and became more conspicuous, after the second operation. In opening her mouth patient herself experienced resistance to the separation of the lips on the left

side. To remedy this defect she consented to another operation, the design of which was to lengthen the mouth on the left side by extending the angle toward the cheek. It was performed June 19th, according to the method described on page 28. Although an abscess formed in the left cheek, adjoining



FIG. 62.

the angle of the mouth, and a small slough was cast off, the final result was satisfactory, and the symmetry of the mouth was restored. In the month of October following, Mr. J. A. Bishop, a skillful dentist of this city, undertook to adapt a plate of vulcanite to the roof of the mouth, that would close the cleft in the bony palate, and also supply four artificial

teeth, that were to take the place of the four upper front incisors, which it was thought best to extract, on account of the irregular position they occupied. Mr. B.'s success was complete. The plate, when finally adapted, was worn with much comfort, and the characteristic defect of articulation, which had existed in a marked degree, was almost entirely corrected. Fig. 62, from a photograph taken in April, 1875, shows the final result.

CASE XXI.—*Congenital Hypertrophy of the Tongue.*

MARY JANE C., aged nine, from Birmingham, Conn., of a healthy family and good constitution, entered St. Luke's Hospital, May 21, 1866, with hypertrophy of the tongue, which, her mother stated, had existed from birth, and had progressively increased with her general growth. The condition of the tongue at the time of her admission was as follows: It protruded habitually from the mouth about two inches beyond the lips, and distended the angles of the mouth, and was overlapped by them. It measured one inch in thickness, two inches and a half in breadth, and four inches and five-eighths in circumference. Its upper surface was coated with a yellowish-brown crust, which on drying became detached and fell off in scales. The protruded portion was of firm consist-

ence and painless. The portion within the line of the teeth, though of full size, was in other respects normal. The under lip hung over upon the chin from the pressure of the tongue. The tongue could be protruded about four inches beyond the front teeth. The lower front teeth were depressed nearly to an horizontal position, and thickly incrustated with tartar,



FIG. 68.

so as to be more than double their normal size. This was true of all the lower incisors and both canine teeth. It was remarkable how little her articulation was affected. She attended school, and was accustomed to recite and sing with her fellow-pupils. The sublingual and submaxillary glands were not enlarged. Her general health and aspect were good. (See Fig. 63.)

First Operation.—Performed May 26th. The head of the patient was supported against the breast of an assistant, who retracted the angles of the mouth with his forefingers. The tongue was then seized with a volsella, and drawn forward out of the mouth. A strong ligature was then passed from below upward, through both edges of the tongue, as far back as possible. The ends of each ligature being tied together, formed loops with which to control the tongue and draw it forward. While held forward and spread out laterally by means of the loops, the tongue was transfixed with a straight, sharp-pointed knife through its middle, from below upward, as far back as the teeth would permit, and a flap formed on the left side by cutting forward and outward. The right flap was formed by applying the knife to the opposite edge of the tongue, and cutting in a reversed direction inward and backward. The arteries spurted briskly, but were readily secured in the usual manner and ligated. The flaps were then brought into contact and secured by sutures. It was now for the first time noticed that the lower jawbone itself had undergone a change of form in front, in consequence of the constant pressure of the tongue. The teeth of the upper and lower jaws came in contact only as far forward as the first molars on both sides, while the front teeth were more than one inch apart at the symphysis.

May 27th.—Patient obtained some sleep; pulse 126; febrile reaction considerable. Ordered tinct. aconit. rad. gutt. j; spirit. Mindereri, ℥ij, q. 4. hor.

28th.—Slept well; pulse 120; swelling of tongue considerable; its extremity has a livid, ashy aspect; breath offensive; right side of the face swollen; deglutition somewhat difficult. Ordered wine, and a mouth-wash of dilute permanganate of potash.

29th.—The lateral flaps have separated, and their confronting surfaces are sloughy.

31st.—Swelling abating; sloughs separating; deglutition improving; general condition satisfactory. Subsequently no considerable loss of substance resulted from the sloughing. Healthy action was gradually reëstablished, and on June 18th cicatrization was complete. Although the excess in the breadth of the tongue had been reduced by the operation, there still remained an excess in its thickness at the end of the stump, which occupied the space between the front teeth, and could not be retracted. A second operation was therefore resorted to, to remedy this defect, and was performed on July 1st following.

Second Operation.—A strong ligature was passed transversely through the back part of the tongue by means of a long darning-needle, and the two ends made use of to hold the tongue forward. The tongue itself was seized with a volsella, applied at opposite points to its edges, near the extremity, and com-

pressed it laterally, so as to increase its thickness. While thus held, a wedge-shaped portion was removed by transfixing the tongue laterally, at a point far back and equidistant from its upper and lower surfaces. The under flap was first formed by cutting forward and downward through its under surface, while the upper flap was formed by applying the edge of the knife upon the upper surface, at a point opposite to the extremity of the newly-formed under flap, and cutting in a reversed direction backward and downward to the point where the first section had been commenced. After ligating two or three arteries, the two flaps were brought into exact contact, and secured by interrupted thread sutures at the margin. By this second operation the tongue was reduced in thickness and length, so that it could now easily be retained within the limits of the dental arch, and its extremity had assumed a rounded, flattened shape. On the second day after the operation the tongue had become much swollen, and considerable febrile reaction had taken place; pulse 124; the surface of the body hot and dry. On the seventh day all the sutures had been removed in succession. There had been no sloughing, nor any separation of the flaps, as after the first operation. The swelling having mostly subsided, deglutition had become easy, and patient was able to be up most of the day. Convalescence progressed favorably till about the 10th of July,

when she began to complain of sore throat, which was accompanied by fever, restlessness, loss of appetite, etc.

July 12th.—The submaxillary glands were hard and tender, and the whole submaxillary region much swollen externally.

16th.—The tongue itself was also much swollen; articulation had become very indistinct and deglutition very difficult; a copious secretion of saliva flowed constantly from the mouth; pulse 144. Ordered poultices of flaxseed-meal to the throat and a Dover's powder at bedtime.

19th.—The swelling has softened and fluctuation is perceptible. Subsequently a spontaneous opening formed under the tongue, and discharged pus abundantly into the mouth, affording great relief, and followed by an abatement of the swelling and a decided amelioration of all the symptoms. Recovery thenceforward progressed rapidly, and without further drawbacks. Before leaving the hospital, Mr. J. A. Bishop removed the tartar incrustation from the teeth, and adapted a fixture with elastic bands, with which to exert upward pressure under the chin, for the purpose of restoring the under jaw to its normal relations to the upper. The fixture was to be worn as much of the time as possible. She returned to her home, in Connecticut, on July 27th. In July of 1869 the author saw the patient, and on examination

found her condition as follows: The tongue retained its normal dimensions, and occupied the floor of the mouth, its upper surface not rising above the level of the lower teeth. When the act of protruding the tongue was attempted, it bowed itself upward and



FIG. 64.

somewhat forward, being sustained by the connections of its under surface. The lower front teeth had regained their erect position, and the separation between the upper and lower teeth in front was scarcely half an inch. Articulation was without any defect. (*See Figs. 64, 65, from photographs taken at that time.*)

Remarks.—The method employed in the second operation that was performed in the above case is obviously the one to be preferred in similar cases of



FIG. 65.

hypertrophy of the tongue. It has the advantage of reducing more effectually the excess of thickness in the organ than the method employed in the first operation, and leaves the tongue of a more natural shape after the operation. It also avoids severing the connections of the muscles inserted into the under surface of the tongue, and, as the incisions do not involve the floor of the mouth, there is no communication opened between the wound and the in-

termuscular spaces in the submaxillary region, and consequently there is less liability to burrowing suppuration.

CASE XXII.—*Congenital Hypertrophy of the Under Lip.*

THOMAS P. A., aged twenty-five, a native of England, a printer by trade, resident in the United States since three years old, was admitted into St. Luke's Hospital, February 5, 1867. From birth his lower lip had been abnormally large, and with it there co-existed a raspberry-colored stain of the surface of the chin and cheeks wherever they were covered by the beard. During youth and early manhood the parts underwent no special change other than an apparent increase of the swelling of the lip and a brighter redness of the neighboring surfaces in hot weather. While exposed to the hardships of military service in the far West, during the late war, a fresh impetus was given to the growth of the lip, and it increased to double its previous size. In August, 1866, successive operations, consisting of the introduction of red-hot needles, were performed by the late Dr. Charles A. Pope, of St. Louis. Sixty insertions in all were made at various intervals within a period of six weeks. Inflammation and slight suppuration followed; the parts became more dis-

tended, and the permanent result, as patient believes, was rather detrimental, the lip remaining larger than before, and retaining a sensation of numbness. He had also less control over its motions than before the operation. Nothing similar to his condition is known to have existed among his relatives. His condition at the time of admission into the hospital was as follows:

The raspberry discoloration occupied the surface on both sides of the face and chin, where the beard grows. The lower lip was more than double the thickness of the upper lip, and proportionately increased in all its other dimensions. It was pendulous, of a soft, flabby consistence, and free from pulsation. In its substance small, hard knots were felt, which had existed only since the insertion of the hot needles. Simultaneous compression of both common carotids had no perceptible effect on the volume of the lip. (*See Fig. 66.*)

Anticipating copious hæmorrhage in the performance of an operation upon the lip, clamps were devised for making compression at both angles of the mouth. They consisted of two flat steel blades, half an inch wide, bent flatwise at right angles, and made to slide lengthwise upon each other on their flattened surfaces by means of a screw. The vertical portion of the blades beyond the angle was two inches in length, and adapted to compress not only the lip

proper, but also the skin below the lip as far down as the edge of the jaw. Before applying the clamp, an incision was made through the mucous membrane on the inside of the lip where it joins the bone, at a point below the angle of the mouth, to allow the end of the distal blade to be forced down in contact



FIG. 66.

with the periosteum as far as the edge of the jaw. The proximal blade was then screwed up, and the included lip and skin below it were tightly compressed between the two blades. Both clamps having been applied in the manner described, an operation was performed on February 27, 1867, as follows:

A letter-V-shaped patch, including about three-fifths of the lip border, equidistant on either side from the angles of the mouth, and having its apex low down in the median line under the chin, was removed. As one of the clamps became accidentally loosened without causing any considerable hæmorrhage, it was evident that the clamps might have been entirely dispensed with. A single artery only, and that at the lower part of the wound, required a ligature. After dividing the mucous membrane along the line of its reflection from the jaw on either side of the wound, the opposite edges of the wound were brought together, and secured in exact coaptation by five pin sutures, inserted at equal distances from each other, below the lip border. Between every two pin sutures a silver-wire suture was added. Three fine-thread sutures were inserted at the vermillion border of the lip, one of them on its buccal surface. No adhesive-plaster was used. In order to destroy the raspberry discoloration of the face, which caused such a conspicuous disfigurement, a disk-shaped cautery-iron, with a smooth, flattened face, heated to redness, was applied to that portion of the surface situated above the angle of the jaw on both sides, and the application was immediately followed by compresses wet in ice-water, and frequently renewed.

The wound did well. On the third day all the

sutures had been removed, and primary union had taken place at all points. Superficial eschars separated from the burnt surfaces, leaving a healthy granulating sore without any remaining raspberry discoloration. After an absence of three weeks from the hospital, patient was readmitted April 2d, to undergo a second operation. The lip, though much improved, still retained its original excess of thickness. The granulation growth upon the cheeks had become too exuberant, and required a single application of solid caustic potassa, after which the nitrate of silver sufficed to regulate it.

A Second Operation was performed April 10th, to reduce the excess in thickness of the under lip, as follows: A prism-shaped strip was excised from the entire length of the lip-border by two parallel incisions, including between them one-third of the thickness of the lip, and penetrating deep into its substance. The edges of the wound were then brought together, and secured by fine-thread sutures, inserted close together. This operation did well, and produced a still further improvement in the lip. Anxious to return home, patient was discharged April 30th. The experimental treatment of the discoloration of the face by the actual cautery had the desired effect of destroying the raspberry tint, but, in consequence of patient's absence during the healing process, the cicatricial surface, resulting from the burn,

was not so smooth and even as it might have been under proper management.

Fig. 67, which shows the patient's present condition, is from a photograph taken in January, 1872, five years after the operations. Dr. I. D. Beebe, of




FIG. 67.

Hamilton, Madison County, N. Y., through whose kind services the photograph was obtained, in a letter to the author, says that, "were it not for the remaining discoloration of the skin, which is considerable, no one would imagine he had had an operation on the lower lip, as the scar shows but very little."

CASE XXIII.—*Abnormal Growth of Hair upon the Forehead.*

F. J., aged thirteen, a native and resident of the city, with fair complexion, light hair, and of a good constitution, has from birth had an abnormal growth of hair upon the left half of the forehead, covering the entire surface between the left eyebrow and the hair of the scalp, above as well as toward the left temple. The hair of the left eyebrow is much coarser than that of the right, while the abnormal growth of hair is short, soft, and fine, like that of a mouse-skin; both are darker colored than the hair of the scalp. The surface of the skin covered by this growth of hair, after being shaved clean, was found to be of a tawny hue, like that of the scrotum, and slightly elevated. (*See Fig. 68.*) Both the patient and his parents being anxious to have this conspicuous disfigurement remedied, its removal was undertaken and successfully accomplished by destroying the abnormal growth, partly by the actual cautery, but chiefly by the application of solid caustic potassa. The first application was made November 15, 1867. A disk-shaped cautery-iron, with smooth, flattened face, was heated to redness and applied over one-fourth of the shaved surface, and held in contact with it until a dark-brownish eschar was produced. Immediately thereafter compresses wet in ice-water were applied

to the part and frequently renewed. By this means patient almost entirely escaped suffering on coming out of the anæsthesia. After the separation of the eschar the sore surface was dressed with simple cerate, and the granulation-growth kept level with the neighboring surface, during the subsequent cicatrization,



FIG. 68.

by the application of solid nitrate of silver, renewed every second or third day. On November 27th a first application of solid caustic potassa was made to another fourth of the surface, by rubbing the end of the stick of caustic into the surface till the skin became disorganized into a soft, pulpy paste. The ac-

tion of the caustic was then neutralized with diluted vinegar. Two subsequent similar applications were required to complete the cauterization of the hairy surface. Nitrate of silver was also used to regulate the granulation-growth till cicatrization was completed. By the end of the following Decem-



FIG. 60.

ber the entire surface had healed. For more than two years afterward the patient remained under the author's observation. The cicatricial surface upon the forehead continued smooth and pliable, and could be moved freely over the underlying parts. It was a little paler than the adjacent skin. No con-

traction affecting the upper eyelid followed. The improvement resulting from the treatment was every way satisfactory. (*See* Fig. 69.)

CASE XXIV.—*Erectile Tumor of Large Size.*

AN infant daughter of Madame de F., six months old, and of good physical development, came under surgical treatment in January, 1871, for an erectile tumor of congenital origin, seated upon the right shoulder, and covering the upper half of the deltoid region and the adjacent acromion process above. Anteriorly it encroached somewhat upon the pectoral region. Its longest diameter was two inches and three-quarters, and its shortest two and three-eighths. The outer half of the tumor was the thickest, and its margin, which was elevated more than one inch above the neighboring skin, was somewhat shelving; the inner half grew thinner and sloping toward its margin, which was but little higher than the adjacent surface. The surface of the tumor was unbroken, and of a bright raspberry-color; no pulsation was perceptible on it, nor were there any enlarged blood-vessels visible on the surrounding surface (Fig. 70). The treatment consisted in the employment of the actual cautery and the solid stick of caustic potassa. A bullet-shaped cautery-iron,

heated to redness, was the instrument used. Patient was anaesthetized with sulphuric ether on each occa-



FIG. 70.

sion of the application of the actual cautery and caustic.

January 14th.—A first application was made. The iron, heated to redness, was sunk into the sub-

stance of the tumor, at one point in the centre and at five or six points at its circumference, and made to char the tissues with which it came into contact. Immediately after the application, compresses wet in ice-water were laid upon the part, and frequently renewed for several hours afterward. The hæmorrhage was slight, and not sufficient to require attention. On emerging from the anæsthesia, patient did not manifest any signs of severe suffering. A moderate degree of febrile reaction supervened on the following day, but soon subsided thereafter; the local inflammation was also moderate. After the separation of the eschars a dressing of simple cerate was employed.

February 11th.—A second application of the actual cautery was made in the same manner as the first.

22d.—A first application of the solid stick of caustic potassa was made by plunging it into the substance of the tumor, and then promptly absorbing the fluids saturated with the caustic, in order to restrict its action within the limits intended; and, in addition to this, vinegar was also freely applied, to more completely neutralize the action of the caustic.

March 8th.—A second application of caustic was made, and on March 22d a third and final application. The suppuration produced by these applications was at no time excessive, and the patient's

general condition was scarcely disturbed during the progress of the treatment. On the 12th of April following, when seen for the last time (in 1871),



FIG. 71.

there were no remains of the erectile tissue, the sore presented a healthy aspect, and was nearly healed.

On December 22, 1875, nearly five years after the treatment, the seat of the tumor was examined, and presented the following appearance: There was

no remaining discoloration, but a pale, smooth cicatricial surface in the place of the tumor. The skin also was pliable, and moved freely on the underlying parts.

Fig. 71, from a photograph taken at this time, shows the condition of the shoulder.

CASE XXV. — *A Pendulous Tumor (Molluscum Fibrosum) arising from the Right Half of the Forehead and Temple.*

Miss R. R., aged thirty-five, a resident of the city, came under surgical treatment on account of a pendulous tumor arising from the right half of the forehead and temple, and involving to some extent the hairy scalp above. The tumor was elevated more than an inch above the level of the surrounding surface, and in the progress of its gradual development had gravitated toward the cheek and become invested with the right eyelids, which were spread out upon its surface. It had also dragged the eyeball itself partly out of the orbit. The tarsal margin of the upper lid, which overlapped and concealed the under lid, formed the lower limit of the tumor, and descended upon the cheek to a point on a level with the middle of the nose. The right eyebrow was spread out upon the surface of the tumor one inch

below the level of the left. The skin covering the tumor retained its natural color, was thinned, and could be gathered into folds between the thumb and fingers. Its substance was of moderately firm consistence, but of loose texture, and the tumor itself glided freely upon the surface underlying it.



FIG. 72.

When grasped in the hand it had almost the pliability of the scrotum. On separating the lids, the eye was found to be natural in appearance, and to possess good vision. (*See Fig. 72.*) When an infant a few months old, her mother first noticed a somewhat elevated ridge across the right half of the forehead, which increased very gradually till it reached its present dimensions. Its growth became accelerated

after her recovery from a dangerous illness in childhood, for the treatment of which she was profusely salivated. When eleven years old, an operation was performed, and a portion of skin of the size of a silver quarter of a dollar excised, but without any benefit. Other growths, varying in size from a small pea to a cranberry, are scattered over the body, upon the limbs and trunk. The patient allowed one of these growths, of a pendulous form, near the fold of the right elbow, to be removed for microscopic examination prior to the performance of an operation on the face. Unfortunately, a memorandum of the result of the examination was lost. It was, however, ascertained that there were no cancerous elements in the structure of the tumor. Patient's general health being favorable, the first operation was performed January 13th.

First Operation.—A transverse incision, commencing above the left eyebrow, was carried across the tumor along the middle of the forehead to the right temple; a second incision, commencing above the right temple, was carried vertically downward to the upper part of the cheek, joining the first incision at its terminus on the temple, the two incisions forming a letter T laid upon its side. The angular flaps of skin thus formed were dissected up from the surface of the tumor to its margin above and below. The tumor itself was then raised at its circumference from the underlying surface, and a prolongation of

it, extending into the orbit under its roof, was dissected out. No large vessels were encountered, nor was the hæmorrhage from small vessels considerable. Ligatures were applied to all vessels requiring it. After the removal of the tumor the redundant skin was pared away, so as to permit a good adjustment of the flaps to each other. Numerous thread sutures were then inserted to secure the adjustment. A good result followed, with primary union at almost all points.

Second Operation.—April 8th. A flattened portion of the tumor, situated high up on the right temple, not having been included in the previous operation, a second operation, of only limited extent, was performed for its removal, and was followed by prompt healing of the wound.

Third Operation.—Performed April 28th. The right eyeball having been attacked with acute inflammation, which proceeded rapidly to disorganization of its internal structures, it was extirpated, and at the same time some remaining portions of the morbid growth were removed. A good result followed the operation, but the remaining conjunctival cavity unfortunately was not sufficiently capacious to permit the insertion of an artificial eye.

Patient was seen as late as September 11, 1873, and her condition ascertained to be as follows: She enjoyed ordinary good health. There had been no

return of the morbid growth; the cicatricial lines upon the forehead and face had become quite indistinct; the right eyelids were habitually closed and somewhat sunken. Fig. 73, from a photograph taken



FIG. 73.

in October, 1873, accurately represents the final result of the three operations.

Remarks.—In vol. xxxvii. of the “*Medico-Chirurgical Transactions of 1854*,” the late Dr. Valentine Mott reported five cases of what he termed “a peculiar form of tumor of the skin,” and denominated “*pachydermatocele*.” Two of the five patients were boys, aged respectively twelve and fourteen years; three were females, aged twenty-four, forty, and forty-five years. In all the five cases the growths were

of congenital origin, or dated from very early childhood; they were all of a pendulous form and lax texture. In the boys the tumors occupied the side of the head and face. In two of the females the side of the chest below the mamma was the seat of the growth, and in the third the tumors occupied the entire left side of the neck, between the larynx and ligamentum nuchæ, and extended upward as high as the ear, and downward upon the sternum, clavicle, shoulder, and scapula; they hung pendulous as low down as the umbilicus. All the patients were operated on for the removal of the tumors, and in all the subsequent healing process advanced kindly. One of the female patients, forty years old, died in a low typhoid state within a few days after the operation, her condition being induced by self-imposed privations of all sorts, from a fancied conception of extreme poverty. Three of the patients, one boy and two females, recovered without any return of the growth; one of these three, a female, forty-five years old, whose case was the most formidable from the number and large size of the tumors, withstood, in the progress of her recovery after an operation, two formidable attacks of erysipelas that endangered her life. One of the four cases of recovery was a boy, fourteen years old, on whom a second operation was performed on account of the return of the growth.

Though both operations did well, the growth reappeared again after the second operation. In regard to the nature of one of the tumors examined, Dr. Mott states, "My colleague, Prof. Lovett, has kindly furnished me with the following notes of the microscopical appearance of one of the tumors: 'The specimen appears to me to consist of an hypertrophy of the skin and the subcutaneous cellular tissue. Under the microscope I find nothing but the exaggeration of the natural tissues; there are no evidences of a malignant formation.'" In vol. lvi. of the "Medico-Chirurgical Transactions," Mr. Pollock, of St. George's Hospital, London, has reported a case of what he terms "molluscum fibrosum, or fibroma." The patient was a widow, thirty-three years old, in whom the first appearance of the growth dated back to her earliest recollection. Besides numerous growths varying in size from a split pea to a small walnut, scattered over different parts of the body, a group of very large-sized tumors arose from the right side of the neck, between the hairy scalp and shoulder behind, and in front between the ear above and a line across the chest four inches below the top of the sternum. They hung in festoons over the right mamma, and reached below the umbilicus. An operation for their removal was followed by a favorable result. Four months after the operation there had been no

return of the growths. Of the structure of these growths Dr. Whipham, after a microscopical examination, reported to Mr. Pollock: "The growth is due partly to an excessive hypertrophy of the connective tissue of the true skin, and partly to an abundant cell-growth occupying interspaces between the bands of fibrous and elastic tissue, which, as has been shown, comprise the chief part of the growth. Neither the large pendulous, nor the smaller sessile tumors, depend upon any alteration of the epidermis, rete-mucosum, glands, or hair-bulbs, as far as can be made out." In vol. xvi. of the "Transactions of the "Pathological Society," London, Dr. H. G. Wight reported the case of an unmarried woman, thirty-four years old, with an enormous group of pendulous tumors arising from the right half of the neck, between the symphysis menti and the spinous processes of the cervical vertebræ. They originated as low down as the fourth dorsal vertebra posteriorly, and anteriorly as low as a line drawn between the two nipples. Below this line they reached to the umbilicus. "The skin," he says, "resembles in some measure that of the scrotum. Patient first noticed the growth when about fourteen years old. She was found dead from suffocation, which, as was supposed, had occurred during an attack of epilepsy, to which she was subject." The structure of the tumors in all the above

cases, including the author's, was strikingly the same, and consisted essentially of an overgrowth of the constituents of the subcutaneous connective tissue.

See also Virchow, "Die krankhaften Geschwülste," vol. i., pp. 350-353.

THIRD CLASS.

CICATRICIAL CONTRACTIONS FOLLOWING BURNS.

THE cases belonging to this class, which have come under the author's own observation, will be arranged in two groups.

First Group.—Comprises lesions involving the face, and producing distortion of the eyelids, nose, and lips. Their treatment calls into requisition the resources of plastic surgery, and consists essentially in first liberating the distorted parts by detaching them from their underlying connections sufficiently to permit them to be restored to their normal relations, and then transplanting sound skin from the nearest available locality, with which to fill up the space made bare by the restoration.

Second Group.—Comprises lesions involving the neck and upper limbs, and producing contractions at their articulations, chiefly in the direction of flexion. In the cases belonging to this group, which are hereinafter narrated, the cicatricial bands producing the contractions consisted almost exclusively of integument, the underlying aponeurosis being involved only

to a slight degree, while the muscles and tendons were not at all concerned. Their treatment consisted—

1. In excising the corded folds of cicatricial integument that maintained the contraction, and then dividing the edges of the resulting wound at every point where any resistance still remained, which prevented complete extension of the part. At some points it was also necessary to dissect up the edges from their underlying connections. In a word, the liberation of the parts required to be so thoroughly performed that they would have the utmost freedom of motion.

2. The liberation of the parts having been accomplished in the manner above described, the next step in the treatment was to adapt a suitable mechanical appliance, that would maintain the parts in their restored position during the process of cicatrization, and for a long time afterward. If the front of the neck should be the part involved, as in Case XXVIII., a brace would be required to keep the chin constantly elevated and the neck upon the stretch. The ingenuity of the surgeon must supply such modifications of the mechanical support as the exigencies of each case may require. It being necessary to wear the support constantly and for a long time, the comfort of the patient should never be lost sight of.

3. A third and very important step of the treatment is, to regulate the process of cicatrization during

its progress and until its final completion, in order that the healing surface may be kept smooth and even. To do this, two expedients are indispensable, and both of them require to be employed with unwearied perseverance. The constant tendency of the granulation-growth to become exuberant makes it necessary to repress it by the frequent application of caustics, of which solid nitrate of silver and caustic potassa have been preferred by the author. In applying the nitrate of silver, it is not sufficient simply to pass the stick over and in contact with the surface of the granulations, as is commonly done. It requires to be buried deeply into their substance at numerous points in close proximity to each other; and often the application has to be repeated at each daily dressing. Even this energetic use of the nitrate of silver will not always be sufficient to control the exuberant growth. It then becomes necessary, with the aid of anæsthesia, to have recourse to the solid caustic potassa, special precautions being taken in using it to restrict its action within the limits intended. This may be done by promptly absorbing with lint the fluids that so readily dissolve the caustic, and become saturated with it; and also by the application of common vinegar, which neutralizes its action. The potassa being much more energetic in its action than the nitrate, is only required at long intervals. In conjunction with the use of caustic,

compression is an efficient and indispensable auxiliary, and is best employed by means of adhesive plaster, cut into strips, and applied in immediate contact with the entire granulating surface. The strips should overlap each other, like shingles on a roof, and they require to be renewed at each daily dressing. If in the progress of cicatrization there should be a tendency to the reproduction of contracting bands or folds, they should be freely divided with the knife at two or more points across the direction of their length. The formation of new skin-growth may further be promoted by leaving small patches or islets of the cicatricial skin at different points upon the raw surface, instead of removing it entirely. These islets will subsequently send forth from their edges in every direction new growth, and thus accelerate the healing process. It should be borne in mind that, for a long time after healing is completed, measures must be perseveringly employed to oppose the tendency to re-contraction, which is very persistent; such measures, for instance, as wearing, during the night and a part of the day, the mechanical support that had been used in the earlier period of the treatment; also, the frequent stretching of the parts by manual force, together with the practice of appropriate gymnastic exercises. The practical application of these several methods of treatment will be best understood from

the description of their use in the subsequent cases. These views of the treatment of cicatricial contractions agree in all important particulars with those taught by Dupuytren in his "Leçons Orales," vol. ii., Art. 1, p. 66, *et seq.*, Paris, 1832. The late Mr. Henry Earle, of London, reported, at a much earlier period than Dupuytren, in the "Medico-Chirurgical Transactions," vols. v. and vii., cases of this injury, involving the neck and upper limbs, successfully treated by the same methods. Mr. G. H. James, Surgeon to Dover and Exeter Hospital, subsequently to Mr. Earle, reported, in vol. xiii. of the "Medico-Chirurgical Transactions," cases successfully treated on the same plan; and in a recent publication "On the Results of Operations for Cicatrices after Burns," London, 1868, Mr. James has given his additional experience.

CASE XXVI.—*Disfigurement of the Face from Cicatricial Contractions; Extirpation of one Eyeball; Closure of the Orbit by a Plastic Operation.*

ANNIE MULLADY, aged ten, of Irish parentage, from Scranton, Pa., was admitted, June 2, 1870, into St. Luke's Hospital. The burns, from the consequences of which her present disfigurement resulted, occurred four years previously in a railroad-

car that was precipitated over an embankment, causing a heated stove to fall upon her. Her condition at the time of her admission was as follows: Both lids of the left eye were entirely destroyed, leaving the eyeball uncovered and projecting beyond the orbit. The eye itself was sightless from dense opacity of the cornea. A condition of extreme irritability kept it in constant motion and affected the sound eye, obliging her habitually to incline the head forward, so as to shun direct exposure to the light. An ulcerated spot, of the size of a silver half-dollar piece, still remained unhealed over the left frontal boss. The entire surface of the left half of the forehead, surrounding the ulcer, was cicatricial, and the same condition extended up over the scalp, which was denuded of hair-growth. The surface of the left cheek and temple, as well as the nose, was in a like cicatricial condition. The nose itself was shrunken, distorted, and drawn over toward the left side to such a degree that the columna nasi no longer corresponded to the inferior border of the septum, but stood off to the left of it, leaving it exposed to view. Both nasal orifices were misshapen and contracted. The upper-lip border, at the left angle of the mouth, was drawn up into a notch. The inner half of the upper lid of the right eye was everted and notched by a cicatricial contraction of the skin above it. The right eye itself had escaped injury,

and possessed good vision. (*See Fig. 74.*) Her general health was good. The first and most urgent indication was to relieve the sound eye of its sym-



FIG. 74.

pathetically-induced irritable condition, by the extirpation of the sightless ball of the left eye.

First Operation was performed June 29th, as follows: The remains of the conjunctiva were detached from the ball of the eye with scissors, and the muscles successively divided at their insertions into the globe of the eye. The optic nerve was then severed at its junction with the eyeball, and the ball itself removed. No arteries required to be ligated.

A covering of soft scraped lint was laid over the part, and secured by a single turn of bandage, lightly applied around the head. Healing progressed gradually, and a slightly-concave surface, bounded by the bony margin of the orbit, remained; the surface consisting, as it did, of conjunctiva, continued to exude a viscid secretion. The effect upon the sound eye of the removal of its fellow was most beneficial. It at once regained a quiescent condition, and was no longer sensitive on exposure to the bright light. The next step to be undertaken was to restore the upper lid of the right eye, which was everted at the inner half of its tarsal margin. For this purpose a

Second Operation was performed, as follows: An incision, commencing above the inner canthus, was carried across the upper lid on a line parallel with the commissure of the lid, to a point upon the right temple above and a little beyond the outer canthus. The everted inner half of the tarsal edge of the upper lid was liberated, and brought down into contact with the opposite tarsal margin of the lower lid, and the edges of the wound were dissected up to afford space for the insertion of new material, which was obtained by raising from the temple a patch of skin, having its pedicle of attachment adjacent to the outer extremity of the bare space that was to be covered, and its long axis extending upward on the

temple. The outlined patch, after being dissected up from its underlying connections, was transferred edgewise (by causing it to perform a circuit of a quadrant) to the space above the upper eyelid, and there accurately adjusted by sutures. The raw surface left upon the temple was covered by approximating the opposite edges of the wound, and securing them in contact also by sutures. Primary union followed, and the notch at the tarsal margin was nearly obliterated. The conjunctival surface covering the left orbit, from which the eyeball had been extirpated, continued to exude a viscid secretion, and presented a conspicuous, unsightly disfigurement. It was therefore decided to conceal it by covering it with a patch of skin, to be taken from the right half of the forehead—that being the nearest point where sound material could be obtained for the purpose. As a preparation for the proposed operation, the secreting conjunctival surface covering the orbit required to be converted into a granulating surface. This was done by the application of a cautery-iron, heated to redness, and was followed by the separation of a superficial eschar, and the substitution of a healthy suppurating surface. After this preparatory step a

Third Operation was undertaken December 22d, as follows: An incision, dividing the skin, was carried around the bony margin of the left orbit, and the outer edge of the incision was dissected up and everted

sufficiently to allow it to be adjusted to the edge of the patch of skin with which the included space was to be covered. To provide material for this purpose, a patch of skin of the requisite size and shape, regulated by a pattern previously cut from oiled-silk, was outlined upon the right half of the forehead by an incision, the long axis of the patch being directed upward, and its pedicle of attachment corresponding below to the inner half of the eyebrow. Its free extremity included a portion of the hairy scalp above. In order to afford a continuous raw surface between the pedicle of the patch and the bare orbital space it was intended to cover, an incision was carried from the edge of the patch at the inner end of the right eyebrow transversely across the forehead to the bare orbital space, and the skin intervening between the patch and space was displaced from above the incision toward the left side, where it retained its attachment, and was reserved for subsequent use. The patch was now brought around edgewise, from right to left, and from above downward, till it reached its destination, and was adjusted by its under surface continuously in contact with the underlying denuded surface prepared for it. The adjustment was secured by pin and thread sutures. The displaced portion of skin, held in reserve, was utilized by transferring it in an opposite direction, from below upward, and from left to right, to the surface left bare by the fore-

head-patch, the lower half of which it filled up, leaving the upper half to be treated with a collodion crust, under which healing might take place by granulation.

The subsequent management and progress require no special notice. Primary union took place at almost all points of the circumference of the patch. A slight discharge of pus, mixed with lachrymal secretion, continued for several weeks to escape from the orbit by a small opening at its outer margin, but at length it dried up spontaneously, and the opening itself finally closed permanently. The collodion-crust became detached and was removed on the tenth day. The underlying surface was in a healthy granulating condition, and thereafter cicatrized progressively till it finally healed. Attempts were next to be made to remedy the remaining deformity of the face, especially the distortion of the nose and upper lip, at the left angle of the mouth. On December 24, 1870, a

Fourth Operation was performed, as follows: An incision, commencing at a point one finger's-breadth below the inner canthus of the right eye, was carried obliquely downward across the nose and left cheek to a point one inch below the left angle of the jaw. In its entire course this incision divided the cicatricial surface. Both edges of the incision, after being dissected up, receded wide apart, so that the nostrils and the left angle of the mouth regained their natural shape. In order to fill up the wound-space thus

produced with new material, a patch of skin of the same shape and size was outlined by an incision upon the left cheek and temple, and dissected up from its underlying connections, but left attached by a broad pedicle below, where it was adjacent to the extremity of the wound-surface which it was intended to cover. The patch itself was now transferred edgewise to its new locality, and the confronted edges of the patch and space were accurately adjusted to each other and secured by sutures. Omitting unessential particulars, it is sufficient to state that the transferred patch sloughed, owing no doubt to the cicatricial condition of its surface. After the sloughs had separated and the surface assumed a healthy granulating condition, another attempt was made to supply new material with which to fill the wound-space. This was attempted by engrafting a patch of skin from the patient's own hand. For this purpose a

Fifth Operation was performed April 14, 1871, as follows: The wound-surface upon the left side of the nose and cheek was first prepared by paring away the granulations with scissors applied flatwise, and trimming the edges afresh after dissecting them up. A patch of skin of the required shape and size was then outlined upon the dorsum of the patient's own left hand, and located with its pedicle resting upon the dorsal surface of the metacarpal bone of the thumb, and its free extremity at the commissure be-

tween the index and middle fingers. The patch was dissected up from its underlying connections, and left attached at its pedicle. The wound-surface on the back of the hand, made bare by raising the patch, was closed by approximating the opposite edges of the wound, and securing them in contact by sutures, before putting the hand in position. The hand itself was now brought up with its palmar surface applied to the left side of the face, and supported in position, while the patch from the hand was transferred to the cheek, and accurately adjusted to the space prepared for it, and there secured by sutures. A succession of assistants was organized to support the hand and arm in position, and maintain them at perfect rest, if possible. At the expiration of forty-eight hours, however, patient could no longer endure the irksomeness of her position, and was relieved by severing the patch at its pedicle. Its changed color had already indicated that its vitality had ceased, and the experiment had failed. The wound upon the dorsum of the hand healed mostly by primary adhesion, and no defect in the functions of the fingers ensued. Not willing to abandon my purpose, another operation was devised, with the view of obtaining sound skin from below the jaw, with which to cover the wound-space upon the cheek. On the 6th of May following a

Sixth Operation was executed, as follows: The wound-surface on the left cheek having again assumed

a healthy granulating condition, it was prepared in the same manner as for the preceding operation. A patch of skin of the requisite shape and size was then taken from the side of the neck under the jaw, the free extremity of the patch extending forward beyond the symphysis under the chin, and its pedicle retaining its connection over the angle of the jaw. After the transfer of the patch to its new locality upon the cheek, and its adjustment by sutures, the wound-surface below the jaw was closed by approximating the opposite edges of the skin, and securing them in contact by sutures. About three-quarters of an inch of the extremity of the patch sloughed within forty-eight hours after its transfer, and primary union failed to take place elsewhere. It therefore became necessary to maintain the remainder of the patch in place during the process of healing by granulation, and especially to hold it well up toward the nose. To accomplish this a beaded-wire clamp-suture was employed in the following manner: A common darning-needle, threaded with flexible silver wire, was passed through the middle of the patch of skin, and then carried through the outer wall of the left nostril, close to the cheek, and into its cavity and out again by its external orifice. A smooth round glass bead was strung upon the end of the wire coming out of the nostril, and the end of the wire itself knotted to hold the bead. The other end

of the wire traversing the patch of skin was also strung with a bead and a perforated leaden shot. By drawing upon this end of the wire, the bead at the opposite end was brought into the cavity of the left nostril, and rested in contact with its outer wall. At the same time the patch of skin was slid up against the side of the nose and cheek, then secured in its place by the bead pressed against it, and held fast by mashing the shot upon the wire with pliers. This contrivance answered a very useful purpose in keeping the patch quietly in place during the process of healing by granulation. The adjustment of the parts was further perfected by compresses kept in place by strips of adhesive plaster, often renewed. The beaded-wire suture was left in place till the twelfth day, when healing was so far advanced that it could be dispensed with. A plug of soft sponge of suitable size was kept in the left nostril to maintain it in good shape, and was renewed daily. On June 9th cicatrization was complete. On the 15th her mother took her home without my knowledge or consent. This abrupt removal from the hospital cut short all further treatment by which a more perfect result might have been attained. On May 19, 1872, I saw patient at her home, in Scranton, and ascertained as follows: There were scarcely any remains of the notch in the upper lip at the left angle of the mouth. The nose was less distorted than before the

treatment. The patch covering the left orbit, consisting as it did of a portion of the hairy scalp, was covered with a growth of hair. This it was intended to destroy had she remained longer in the hospital.




FIG. 75.

Fig. 75, showing the final result, was from a photograph taken at a gallery in Scranton, at my request.

CASE XXVII.—*Cicatricial Contractions involving the Chin and Front of the Neck.*

ALEXANDER S., residing with his parents in the upper part of this city, was, in February, 1869, at the age of two years and one month, severely burned from his clothes taking fire. The chin and front of the neck were the parts most seriously involved. In the month of June following, the burnt surface having healed, patient accompanied his family to their residence in the country. In October, 1870, when he came under the author's care for surgical treatment, being at the time three years and nine months old, his condition was as follows: A broad band of cicatricial skin of dense structure occupied the front of the neck, extending in a direct line downward, from the lower border of the under jaw to the top of the sternum and clavicles, and approximating the parts so that the jaw could not be elevated beyond four inches above the sternum. The connections of the band with the jaw laterally reached from the left angle of the jaw to a point below the right angle of the mouth. Its connections below with the top of the sternum and upper surface of the clavicles corresponded in extent laterally with those above. The prominence of the chin and the profile outline of the front of the neck were both obliterated. The separation of the jaws and lips from each other left

the tongue habitually exposed to view, and allowed the saliva to dribble constantly from the mouth. The anterior surface of the band was uneven, and traversed in a vertical direction by alternate ridges and furrows. From both edges of the band sound skin extended upon the sides of the neck, after first receding behind the band and forming there a pocket of such a depth that the fingers, when pressed toward each other from opposite directions behind the band, could be made to meet together in the median line, with the skin only intervening between them. When the band was put upon the stretch by elevating the head, the left half of the under lip was drawn down more than the right. Patient also suffered from a tormenting itching, that would suddenly attack the cicatricial parts, and provoke him to scratch and pull upon the band in an almost frantic manner. His mother was of the opinion that for several months past the condition of the parts had undergone no change. His general health was good. (*See Fig. 76.*)

First Operation.—Performed at patient's residence, November 5, 1870, with the aid of Prof. A. C. Post and Drs. C. M. Bell, J. N. Beekman, and Robert Watts, as follows: The entire cicatricial band was divided into three serrated angular flaps by two diverging incisions, carried from the symphysis menti downward and outward to either lateral margin of

the band, where it joined the clavicles. From these terminal points incisions were made, one along either margin of the band, upward and outward to the lower edge of the jaw. The three flaps thus formed were then dissected up from the subjacent loose connective tissue, beginning at their apices, and proceeding toward their bases, at which latter points the dissection was carried a short distance beyond



FIG. 76.

the limits of the cicatricial skin. The head could now be thrown back and rotated freely in every direction. The surface thus laid bare involved the entire anterior region of the neck. The next step was to reapply and adapt the detached flaps to this extensive denuded surface, while the head was kept in an elevated position. In effecting this adjustment it was necessary to excise redundant folds and pare off the edges of the flaps, in order to adapt them to

each other. On the left side of the neck it was also found necessary, in order to relieve tension and facilitate the approximation of the edges of the flaps, to make an incision, five inches long, through the thickness of the skin across the base of the neck, on a line parallel with the edge of the wound and two inches distant from it. The edges of this incision gaped widely apart, and thereby the desired end was accomplished. The raw surface, resulting from the side incision, was coated over with collodion-crust and left to heal by granulation. A covering of scraped lint was spread over the other parts, and secured by strips of adhesive-plaster. Though the operation was necessarily protracted, and the loss of blood considerable, there was no extreme depression of the vital powers in consequence. A good degree of reaction followed, and, aided by an anodyne, the patient passed the first night pretty comfortably. At the expiration of forty-eight hours sloughing had destroyed all the flaps, except a portion about one inch in breadth along their bases. The sutures were removed in succession, and the sloughs got rid of as fast as they separated. Special care was taken, in dressing the wound, to hold the detached skin, that had escaped sloughing, in close contact with the subjacent surface by means of long strips of adhesive-plaster, carried high up on either side of the face and over the temples, and on the sides of the neck. At

the same time the head was also kept well elevated. The wound took on healthy action after the separation of the sloughs, and the patient's general condition was all that could be desired. The detached skin became adherent, and cicatrization progressed favorably. The exuberant growth of granulations was repressed by the energetic use of solid nitrate of silver, not merely passed over the surface, but plunged deep into the substance of the granulations. This was sometimes repeated daily. Solid caustic potassa was also applied, but at much longer intervals, and only at points where the growth was not sufficiently controlled by the nitrate of silver. At the expiration of about four weeks, when the dimensions of the sore had considerably diminished, a stiff leather stock, protected by a covering of canton flannel, was adapted to the neck and worn constantly, so as to keep the head elevated and oppose the disposition to recontraction in the direction of flexion. As cicatrization advanced, the newly-formed cicatricial tissue manifested a tendency to form salient corded bands, which, if left uncontrolled, would have reproduced to a greater or less degree the original deformity. To prevent this effect, the bands were divided at two or more points across their entire thickness, and to a short distance on either side through the neighboring skin, and deep enough to expose the subjacent loose connective tissue, thus

permitting the fresh-cut edges to retract widely apart. This had the effect of breaking up the continuity of the bands and neutralizing their action. These operations were repeated successively, with the aid of



FIG. 77.

etherization, on the 7th, 12th, and 29th of December. The leather stock, worn for the support of the head, proved after a time objectionable, on account of its chafing the skin and producing ulceration. While

endeavoring to devise some substitute for the stock, my attention was directed to a brace, used by Drs. W. E. Vermilye and C. T. Poore, in the treatment of caries of the cervical vertebræ, which seemed admirably suited to my purpose. At my request, Dr. Poore adapted one to my patient. On the 21st of January it was applied, and has been worn constantly since, except at night. It consists (*see* Fig. 77) of two padded steel bands, arranged parallel to each other,




FIG. 78.

one on either side of the spine, and adapted flatwise to the natural curve of the back. It extends lengthwise from the last cervical vertebra to the top of the sacrum.

These vertical bands are joined below by a broad padded metallic band, which passes half round the body behind and just above the hips. At their upper ends the two vertical bands are joined by a cross-piece, *c* (Fig. 78), to which a steel ring or collar, *d*, of an oval shape, is joined by an upright piece, *c d*, in such a manner as to stand horizontally, and afford a support in front to the chin. A segment of the ring in front, *e*, where it corresponds to the chin, is covered with

chamois-leather, and forms a shelf for the chin to rest upon. On one side, near its middle, the ring has a hinge-joint, *f*, which permits it to be opened in two halves, and hence facilitates its removal and re-application. By means of a screw at the joint over



FIG. 79.

the nape of the neck, a lever action is made to elevate the ring in front and regulate the height of the chin. Two shoulder-straps, *gg* (Fig. 77), and an apron, *h*, with three straps at either lateral edge, serve to fasten the brace in close contact with the body.

The brace, besides supporting the chin in an ele-

vated position, and thereby resisting the recontraction of the cicatricial formation in a vertical direction, exerts, by means of the straps which pass over the front of the shoulders, a constant outward traction upon the skin covering the lower part of the neck and upper part of the chest, the effect of which is also to resist contraction laterally and keep the cicatricial surface flat and smooth. Another important advantage of the brace is, that it compels the patient, whenever he wishes to move his head in any direction, to elevate it so as to clear the chin-piece, or, in other words, by voluntary muscular action to stretch the cicatricial surface in front. In addition to the brace, patient has worn constantly, night and day, a cravat of canton flannel, two fingers wide, secured around and in close contact with the neck by an elastic strap and buckle, for the purpose of holding the new cicatricial surface in contact with the subjacent parts. From its first application the patient has worn the brace uninterruptedly, except at night, and so comfortably as scarcely to restrict his activity or enjoyment. A progressive improvement has taken place in the under lip. It has regained its natural form and position, and the saliva no longer escapes uncontrolled.

Fig. 79, copied from a photograph taken April 22d, shows the final permanent result. The contour of the chin and front of the neck were restored to

their natural form and dimensions, and the head enjoyed entire freedom of motion in all directions. The newly-healed surface presented cicatricial lines radiating in different directions, downward as low as the second rib, upward as high as the left angle of the jaw, and also on both sides of the neck. The cicatricial parts were mostly smooth and level, and pliable upon their underlying surfaces. During the hot weather of August, 1871, the brace was left off entirely for a time, and without any detriment. Subsequently, by way of precaution, it was resumed and worn two days in the week for a while longer, and then dispensed with altogether. On January 1, 1874, patient was examined, and a further improvement found to have taken place. The skin covering the front of the neck and chin had become more supple and pliable. The under lip still maintained its normal relations to the upper lip, and performed all its functions, and the head enjoyed the utmost freedom of motion in all directions.

CASE XXVIII.—*Cicatricial Contractions involving the Face and Hand.*

H. C. W., aged four years and seven months, a resident of New Jersey, of sound constitution and enjoying good health, was extensively burned, when

sixteen months old, by overturning upon himself a kerosene study-lamp. At the expiration of about fourteen months the burnt surfaces had mostly healed. The parts involved in the cicatricial contractions which resulted from the burns were, the left half of the face and scalp and the dorsal surface of the left hand and forearm. When I first examined him, in February, 1872, his condition was as follows: The left half of the scalp, as far back as the occipital region, was bare, and the surface presented a pale, shining, cicatricial aspect; the skin, however, was pliable, and moved freely on the underlying parts. The left ear, though diminished in size by the loss of its rim, retained its natural shape, and was not adherent to the scalp. The skin covering the left half of the forehead and temple, though cicatricial on its surface, was still movable on the subjacent parts. The outer half of the left upper eyelid was everted to an extreme degree, and spread out upon the eyebrow; the inner half of its tarsal border alone came into contact with the lower lid when they were closed. The conjunctiva covering the everted portion of the lid being relaxed and swollen, filled up the space between the lids when closed, and thus protected the cornea, which otherwise would have become opaque from constant exposure. The eyelashes, as well as the tarsal edges of both lids, had escaped injury. The eyebrow was denuded of hair. The

eyeball itself had sustained no damage, the cornea retained its natural lustre, and vision was unimpaired. The surface of the left cheek and side of the nose was cicatricial, and the contraction consequent thereupon had drawn the under lid away from contact with the eyeball, but without producing any eversion of its tarsal margin. The left angle of the mouth was somewhat drawn upward by a vertical fold of cicatricial skin upon the cheek, immediately above it. In consequence of the condition of the lids of the left eye, patient habitually held his head inclined forward and to the left side, so as to avoid direct exposure to bright light, and had thereby acquired a peculiar expression of countenance (*see* Fig. 80). A description of the left hand and forearm will be given in a subsequent part of the narrative.

On the 6th of February, 1872, an operation was performed in the presence of Dr. James A. Davis, the family physician, from Bloomfield, N. J., Drs. John N. Beekman and Thomas E. Satterthwaite, and Prof. A. C. Post, M. D.

First Operation.—The object of this operation was, to restore the upper lid of the left eye to its normal relations and functions. It was attempted as follows: Two incisions were started from a single point high up on the forehead, above the middle of the left eyebrow, and continued downward in lines diverging from each other and terminating, one at

either canthus of the eye. The inverted V-shaped patch of skin, included between these incisions, was dissected up from the pericranium as low down as the bony margin of the orbit. The upper lid was thus liberated and brought down so as to allow its tarsal margin to be adjusted in contact with that of the under lid. Before proceeding further, a trans-



FIG. 80.

verse fold of the redundant conjunctiva, lining the upper lid, was excised as far back from its tarsal margin as possible. With the descent of the upper lid, the inverted V-shaped patch of skin, which had been raised from the forehead, was brought down

and secured at a lower level by a pin suture, inserted on either side, together with additional thread sutures above the pin sutures. The surface left bare on the forehead by the descent of the patch was closed by approximating the opposite edges of the wound, and securing them in contact by thread sutures. To relieve these last sutures from tension, parallel incisions were made through the skin on either side, the one over the left temple being three inches, while the other, toward the middle of the forehead, was only one inch in length. The edges of these incisions gaped widely apart, and the desired result was thus obtained. The bare surfaces left were coated with a collodion-crust. In order to hold the tarsal edges of the lids more exactly in coaptation, a beaded silver-wire clamp-suture (*see* page 17) was passed in a vertical direction through both lids, near the outer canthus, and out of the way of the cornea, and left *in situ* for three days. Wet dressings were avoided, and a layer of woven lint of double thickness was kept applied to the parts. Moderate febrile reaction followed the operation, but it subsided on the third day. On the fourth day the last sutures were removed. Sloughing, however, had already taken place, and had involved about three-fifths of the transplanted patch. Healthy suppuration succeeded the separation of the sloughs, and the ulcerated surface progressively di-

minished in size. The collodion-crust separated from the forehead on the sixth day, and from the left temple on the tenth, leaving healthy granulating surfaces to heal by cicatrization. The eyeball had in no way suffered from the presence of the silver wire which traversed the lids. Patient rapidly regained his spirits, and, at the end of one week, resumed his accustomed amusements, enjoyed his meals, and rested well at night. The sloughing which had taken place was no doubt to be attributed to the cicatricial condition of the transplanted patch of skin. The ulcerated surface above the left eyebrow progressively diminished in size, and, at the end of the fourth week, measured one inch in its transverse diameter, by three-fourths of an inch vertically. If left to itself it was feared that the contraction consequent upon cicatrization might reproduce, to a greater or less degree, the original eversion of the upper lid. To prevent this it was determined to transfer a portion of sound skin from the right half of the forehead and engraft it upon the ulcerated surface. The operation for accomplishing this object was performed on the 8th of March.

Second Operation.—The ulcerated spot itself was prepared by first excising the granulating surface down to the level of the pericranium, with scissors, applied flatwise, and then paring afresh the edges and everting them slightly. A transverse incision was

then carried across the forehead, on a line continuous with the lower margin of the spot just prepared, and as far as the inner extremity of the right eyebrow. The upper edge of this incision was dissected up to afford a bare surface of one finger's-breadth, which would be continuous with and form a part of the space above the eyebrow. A pattern, cut from oiled silk, of the size and shape of the space just prepared, was applied upon the right half of the forehead, in a vertical position, with its base resting upon the inner half of the eyebrow, and its free extremity involving the hairy scalp above, which had been previously shaved clean. An incision was then carried around the margin of the pattern, and the included underlying patch of skin was dissected up from the pericranium, but left connected, for support, at the margin of the orbit. The pedicle of the patch, at its inner edge, toward the median line, was adjacent to the bare surface which it was intended to cover. Additional room had to be made for the transfer of the patch, by dissecting up the skin from the forehead, above the nose, and displacing it toward the left side, where it remained attached, and was reserved for subsequent use. The patch was now brought down edgewise from right to left, and adjusted accurately to the edges of the space prepared for it, with sutures inserted close together. In order to utilize the portion of skin displaced from above

the nose, for the purpose of covering the surface left bare on the right half of the forehead, it was carried upward from left to right and adjusted to the lower part of the bare surface by means of sutures. The remaining upper portion of the bare surface was coated over with a collodion-crust, and left to heal by granulation. Before proceeding to the operation just described, a transverse fold was excised, for the second time, from the still redundant conjunctival lining of the upper lid, and the lids themselves were secured together by a single-thread suture inserted through the skin alone, near their tarsal edges, and toward the outer canthus. A strip of woven lint, saturated with collodion, was applied across the outer half of the closed lids, to afford additional support. The entire operation occupied one hour and a half, and was well borne, the loss of blood having been inconsiderable. A layer of woven lint, of double thickness, spread with cerate to prevent its sticking to the surface, was laid upon the forehead. Elixir opii, gutt. x, and weak brandy-and-water were prescribed.

March 10th.—Second Day.—Slight febrile reaction, and moderate inflammatory tumefaction. Removed three pins, and changed the yarn on the remaining ones.

12th.—Fourth Day.—All the sutures have been removed in succession, including the one holding the

lids in contact. To supply the place of this last suture strips of adhesive-plaster were applied. Primary union has taken place at almost all points, and without any sloughing.

20th.—*Twelfth Day.*—The collodion-crust came off from the forehead, exposing healthy granulations at the circumference of the sore, but in the centre a brownish patch of sloughing pericranium, which had been unintentionally divided in the operation. No exfoliation of bone, however, followed the sloughing of the pericranium; healthy granulations covered the spot, and cicatrization followed. Iron and quinine were ordered as a tonic.

Third Operation.—The patient being in excellent condition, from a stay of three weeks in the country, a third operation was performed on the 20th of April, for the purpose of removing a conspicuous distortion of the left angle of the mouth. An incision, commencing at a point on the left cheek, bordering on the middle of the nose, was carried downward and outward across the cheek to a point a little anterior to the angle of the jaw. In its course the incision divided the cicatricial fold which drew up the angle of the mouth, and so allowed the latter to regain its natural shape. The edges of the incision, after having been dissected up, receded from each other and left a space between them of about one finger's-breadth. To fill this up with sound skin,

the following method was adopted: A patch of skin of the required shape and size, with its pedicle of attachment adjoining the space to be filled up, and its free extremity on a line below the symphysis, was dissected up from the side of the neck under the jaw. This patch was then transferred edgewise to the space prepared for it, and there accurately adjusted by sutures. The wound left on the neck was closed by approximating its opposite edges, and securing them together with sutures. The treatment was the same as after the previous operations. Union failed to take place, and sloughing of about three-fifths of the patch followed.

On the fourth day the slough had separated, and healthy suppuration succeeded. It was now important to prevent any shrinking of what remained of the patch, and to maintain it in place. This was done by adhesive straps carefully adapted and frequently renewed.

April 25th.—A mild attack of erysipelas developed itself upon the left ear and neighboring surface of the scalp, but soon passed off without any serious consequences. From this time his general health improved, and cicatrization of all the sore surfaces progressed steadily till June 8th, when all had finally healed. The result of the last operation, notwithstanding the loss of so large a portion of the transplanted patch of skin, was a certain degree of

improvement of the angle of the mouth. As no further operations could well be undertaken till the autumn, patient's nurse was instructed to manipulate the parts upon the forehead and left cheek daily, so as to increase their pliability and prevent contraction from taking place. The good effect of these manipulations was manifest when patient returned in October to spend the winter in the city with his family.

The distortion of the left hand, the treatment of which was now to be undertaken, was the result of the same burn that had disfigured the left half of the face. The condition of the hand was as follows: The thumb and all the fingers, though flexed at their phalangeal articulations, were drawn backward toward the dorsum of the hand to such a degree that the proximal ends of the first phalangeal bones rested on the dorsal surfaces of their supporting metacarpal bones. This distorted condition, which existed to a greater degree in the thumb and index than in the other fingers, was maintained by salient folds, consisting of cicatricial skin alone, that were given off from the dorsum of the wrist in radiating lines, one fold proceeding to the thumb and each finger. While the middle, ring, and little fingers retained their parallel position toward each other, the index was widely abducted from the middle finger, and the thumb from the index-finger.

The hand, notwithstanding its distorted condition,

still performed useful service. Flexion and extension at the elbow-joint, as well as pronation and supination of the forearm, remained unimpaired. (*See* Fig. 81.)




FIG. 81.

Preparatory to an operation, a guttered splint, made of tin, was adapted to the palmar surface of the forearm and lengthened out at the wrist by the addition of a flat piece, bent flatwise in the direction of flexion, and adapted for the support of the hand. On the 24th of October, 1872, a fourth operation was performed.

Fourth Operation.—The second, third, and fourth

fingers being similarly involved, their treatment only was to be attempted in the present operation, which was executed as follows: By flexing the fingers and putting on the stretch the cicatricial folds on the dorsum of the hand, a broad longitudinal fold of skin was also rendered tense and salient upon the forearm above. This fold was transfixed at its base, near the middle of the forearm, while gathered up between the thumb and fingers, and a tongue-shaped flap of skin was formed by cutting downward toward the wrist and outward through the surface. The detached flap receded toward the elbow, and the three fingers could now be flexed to a right angle at their metacarpo-phalangeal articulations; the phalanges themselves, however, could not be flexed upon each other, owing to the remaining resistance of the edges of the wound just made upon the forearm, which became very tense and unyielding whenever flexion was attempted. These edges were therefore freely divided across at selected points till there remained no longer any resistance, and complete flexion of all the phalanges could be effected. No tendons or muscles were exposed; the surface laid bare consisted of adipose and connective tissue. A single vessel only required to be ligated. Here and there a suture was inserted to hold the detached skin in place, but without any attempt to procure adhesion. The limb was then placed upon the splint prepared

for it, and the three fingers were secured in a flexed position, by adhesive-plaster, to the flat piece that joined the splint at an angle of flexion, opposite to the wrist-joint. The raw surface, which involved the dorsum of the hand and lower two-thirds of the forearm, was coated with collodion-crust.

After completing this operation, advantage was taken of the anæsthetic condition of the patient to perform another operation for the further improvement of the lids of the left eye. The outer half of the tarsal edge of the upper lid was still disposed to become everted, and this tendency was promoted by the presence of a mass of granulations in the conjunctival *cul-de-sac* at the outer canthus. The commissure of the lids had also become lengthened in a marked degree, so as to exceed the length of the commissure of the lids of the other eye by about one-quarter of an inch. The mass of granulations was first excised by seizing it with a fine-clawed forceps and clipping it off at its base with a scissors curved flat-wise. The attempt was next made to bring about permanent adhesion between the tarsal edges of the lids at the outer canthus. While the lids were held wide apart, and the eye itself was protected by a wet rag, a ball-shaped cautery-iron of the size of a pea, heated to redness, was buried in the *cul-de-sac* of the conjunctival cavity, at the outer canthus, and applied thoroughly to both tarsal edges of the lids

for a distance of nearly two lines from their junction with each other. Compresses wet with ice-water were immediately applied, and afterward frequently renewed.

On coming out from the anæsthesia, patient did not appear to suffer from this severe application. No febrile reaction followed. He passed a good night without an anodyne, and the next day sat up with his arm supported in a sling. Scarcely any redness or swelling of the hand or forehead supervened.

On the fifth day, October 29th, the collodion-crust came off from the wrist and forearm, leaving a healthy granulating surface, which thereafter was dressed with simple cerate, the limb being kept upon the splint. A moderate degree of inflammation followed the application of the actual cautery to the conjunctiva and tarsus. No injury was sustained by the cornea or ball of the eye. Superficial eschars separated, and healthy suppuration followed. Upon the forearm portions of cicatricial skin, which had been detached from their underlying connections, sloughed.

November 5th.—The exuberant granulation-growth has required the energetic application of solid nitrate of silver at every daily dressing. Forcible flexion of the fingers and wrist has been daily practised: this, though a severe process during its performance, was not followed by any lasting pain afterward.

It now became apparent that a great advantage would have been gained by liberating the index-finger, and restoring it to its normal relations at the same time with the other three fingers; it was accordingly decided to do it without further delay. The operation was performed on the 5th of November.

Fifth Operation.—Two incisions, beginning one at the commissure between the thumb and index-finger, the other between the index and middle fingers, were carried through the skin, upward, in converging lines, until they met at a point above the cicatricial fold of skin which held the finger extended. After these incisions had been made, the index-finger could be flexed at its metacarpo-phalangeal articulation, but not at its phalangeal joints. By dissecting up the skin on the dorsal surface of the first phalanx from its underlying connections, the power of flexing the finger at these articulations was fully restored. The index-finger was then brought down by the side of its fellow, where it was secured, and thereafter treated in common with them.

The cauterized surfaces at the outer canthus of the left eye being now in a state of healthy suppuration, and all swelling having subsided, the tarsal edges of the lids were secured in exact contact by a beaded silver-wire suture (*see* page 17) inserted in

a vertical direction through both lids, at a distance of half an inch from their tarsal edges, and at the same distance from the outer canthus. Two fine-thread sutures were also inserted at the edges, after first scraping them with a dull-edged knife. The thread sutures were removed on the third, and the



FIG. 82.

wire suture on the eighth day. Permanent adhesion was thus secured between the tarsal edges for a distance of nearly three lines from the outer canthus, thus shortening the commissure so as to correspond with that of the other eye.

Early in the month of January, 1873, the wound-

surfaces on the dorsum of the hand and wrist had healed. The manipulation of the wrist and finger joints had been continued daily with decided benefit, and the limb had been kept constantly secured to the splint. The thumb, not having yet been subjected to any treatment, remained unchanged. To remedy this deformity was the object of the

Sixth Operation, *February 12, 1873.*—By stretching the thumb in the direction of flexion, a longitudinal fold of cicatricial skin was brought into prominence upon the radial border of the wrist and forearm. At first broad and embracing the root of the thumb, this fold grew narrow as it extended obliquely upward on the palmar surface of the forearm, in the middle of which it became corded, and continued so to the elbow. This fold in the process of its formation had evidently drawn up the thumb into its distorted position, and was still the chief obstacle in the way of its performing complete flexion. This fold was divided, while held on the stretch, by a transverse incision passing one-third around the wrist, and at a distance of one inch above the wrist-joint. The tissue creaked under the knife, and the edges of the wound gaped widely apart, but without affording much relief to the thumb. The subjacent aponeurotic layer was also found tense and resisting, and had to be divided across. Some degree of flexion was thus obtained. It was now as-

certained that the proximal end of the first phalanx was dislocated, and rested on the dorsal surface of the end of its supporting metacarpal bone. In order still further to liberate the thumb, and reduce the dislocation, a longitudinal incision of the skin was



FIG. 83.

carried between the metacarpi of the thumb and the index-finger upward to join the incision across the wrist, and the skin was dissected up on the dorsal surface of the metacarpus of the thumb. All resistance was at length removed, and the thumb could

now be brought down and freely flexed at all its joints. To the splint that had served hitherto for the treatment of the fingers, an addition was made of a separate digitation for the support of the thumb, which required to be maintained in an abducted as well as a flexed position. The newly-made raw surfaces were treated in the usual manner. Patient did well after this, as he had done after the previous operations. The same process of forcible flexion and stretching was employed upon the thumb, as had been used upon the fingers with such good results. In order to reduce the size of the thumb, and improve its shape, it was kept tightly wound with strips of adhesive-plaster. The commissure between the thumb and index-finger was also kept crowded up toward the metacarpus by strips of adhesive-plaster, applied tightly over a saddle-shaped compress made with several thicknesses of woven lint.

On the 1st of May a spot on the wrist, at the root of the thumb, of the size of a finger-nail, was all that still remained unhealed. Under the constant use of the manipulations already described, and the wearing of the splint, the thumb and fingers had regained their natural shape and freedom of motion, and were gradually recovering their power to grasp objects. The thumb, however, in consequence of its greater degree of distortion originally, had not yet

recovered the power of flexion so perfectly as the fingers.

The face had also much improved; the commissures of the eyelids were alike in length on both sides, and the tarsal edges of both lids of the left



FIG. 84.

eye were perfect in their adjustment to each other. A spot upon the forehead, above the left eyebrow, of the size of a thumb-nail, was covered with a growth of hair that required to be shaved two or three times a week. It was a portion of the hairy

scalp, and formed the extremity of the patch of skin which had been transplanted from the right to the left half of the forehead.

On the 10th of May patient accompanied his family on a visit to Europe. From that time to the present (January, 1874), the daily manipulation of the fingers has been kept up regularly, and the splint has been worn at night only, the hand being left free in the daytime to be exercised in every possible way. The father, in a recent letter, says: "His hand has improved, and he now uses the thumb quite a good deal; in fact, for all practical purposes, it is about as useful as the other. We still keep it in the splint at night, and continue the manipulations daily on the hand and also on the face."

Figs. 82, 83, 84, showing the result of treatment, are from photographs taken in Florence, Italy, in January, 1874.

Patient was seen January 15, 1876, and the following particulars ascertained: The lids of the left eye maintain their normal relations to each other, and perform their functions perfectly. A marked improvement of the entire left half of the face is noticeable. The fingers of the left hand perform all their movements to the fullest extent; the thumb only does not admit of flexion to the same degree as the thumb of the other hand, owing to the incomplete reduction of the luxation that had so long previously ex-

isted at its metacarpo-phalangeal articulation. This defect does not, however, in any way impair the conjoint use of the fingers and thumb. As a proof of the complete restoration of their functions, he is now taking lessons on the piano, and his teacher regards him as one of his most proficient pupils.

CASE XXIX.—*Cicatricial Contractions involving the Right Axilla and Arm.*

DANIEL CARRIGAN, aged six years, of Irish parentage, was admitted into the Presbyterian Hospital March 17, 1874. Four years previously an extensive burn, caused by his clothes taking fire, produced the condition for the relief of which he entered the hospital, and which was as follows: With the right arm elevated to an horizontal position, a fan-shaped fold of sound (non-cicatricial) skin, attenuated to such a degree as to be translucent in the sunlight, was developed between the arm and thorax, with its free border stretching from the elbow to the ninth rib, where it terminated in three small radiating folds. This extensive fold involved and constituted the anterior fold of the axilla. Between it and the posterior fold of the axilla, which did not much exceed its normal dimensions, a deep hollow.

extended high up under the shoulder-joint. The skin covering the anterior surface of the arm and forearm (including the wrist), and of the deltoid re-



FIG. 85.

gion above, was cicatricial and of unequal thickness, but still pliable and movable on the underlying surface. At the inner margin of the elbow, where the axillary fold terminated, it was continuous with a

thickened band of cicatricial skin, which stood out in relief upon the forearm when the limb was put upon the stretch by raising it to an horizontal position. This band extended obliquely down over the forearm to the radial margin of the wrist, and prevented complete extension of the elbow and wrist joints, as well as the further elevation of the arm itself, and it also restricted pronation of the forearm. The shoulder, elbow, and wrist joints, were all free. (*See* Fig. 85.)

First Operation.— This was performed March 26th, as follows: The axillary fold of the skin, being put upon the stretch by elevating the arm, was transfixed with a sharp-pointed straight knife at a point high up under the shoulder, and the fold severed from the arm by an incision continued along the line of their junction, and brought out at a short distance above the elbow. The two thicknesses of skin constituting the fold receded from each other, and spread out over the thorax. The cicatricial band, running lengthwise upon the anterior surface of the forearm as a prolongation of the axillary fold, was next pared away with curved scissors applied flatwise along its base. The arm could now be elevated somewhat higher but not to its fullest extent, owing to the resistance still existing at the edges of the wound just made upon the arm and forearm. These, when further elevation of the arm was attempted, became

very tense and unyielding. They were therefore divided across at different points, and also dissected up from their underlying connections until at length all resistance was overcome, and the arm could be elevated to a vertical position, with the elbow and wrist simultaneously straightened out. No muscles or tendons were divided or laid bare; the exposed raw surface consisted of aponeurosis, adipose, and connective tissue. A small islet of cicatricial skin, of the size of a split-pea, was left isolated upon the raw surface at a point near the fold of the elbow, in the hope that it might serve as a point of departure for new growth in the subsequent process of cicatrization. A single small artery only required to be ligated. After hæmorrhage had entirely ceased, the raw surface was covered with a collodion-crust. Patient was then transferred to his bed, and the arm secured in an extended and elevated position by means of a silk handkerchief, rolled diagonally into a cord, which was attached to the wrist and then fastened to one of the crossbars at the head of the bedstead.

27th.—Slept comfortably without an anodyne. Face flushed, moderate febrile heat, and acceleration of the pulse.

28th.—Another good night's sleep; febrile reaction has abated; patient tractable and submissive. Discharge from under the collodion-crust offensive;

removed it entire (on the sixth day), and found the underlying surface of a florid, healthy aspect. Applied simple dressing, with compress and roller-bandage. The arm is kept constantly elevated, and in an extended position.

April 6th.—Commenced dressing the sore surfaces with strips of adhesive-plaster, applied in immediate contact with the surface, the strips overlapping each other like shingles on a roof.

This dressing, which requires to be renewed daily, exerts equable compression over the granulating surface, and represses exuberant growth. It does not, however, suffice alone, but the application of solid nitrate of silver has to be conjoined with it at all points where the growth becomes too luxuriant.

June 10th.—Cicatrization has steadily progressed since the preceding report, and is now nearly completed. On two occasions it has been necessary to resort to the application of solid caustic potassa, the nitrate of silver failing to act with sufficient repressive energy.

A straight, guttered splint, adapted to the dorsal aspect of the entire arm and forearm, has been constantly worn for the purpose of maintaining complete extension at the elbow-joint. In order that the patient might be out of bed, and have the benefit of exercise out-of-doors, a fixture was adapted to maintain the arm constantly in an extended and

elevated position. It consisted of a broad belt of webbing, secured around the body by straps and buckles in front. Upon the belt, where it crossed the back, a socket of stiff leather was fastened in an upright position, and, in this socket, a wooden rod was supported erect, with its upper end reaching above the top of the patient's head. The wrist was secured to the upper end of the rod by a silk handkerchief, rolled diagonally into a cord, and the arm by this means was maintained constantly in an elevated position.

July 15th.—Cicatrization being now complete, the wearing of the fixtures was discontinued and free use of the arm allowed. Certain gymnastic exercises were, however, to be daily practised, which would tend to stretch and elongate the newly-cicatrized parts in the axilla and about the elbow. Though these exercises were perseveringly practised for five months, and had increased the suppleness and mobility of the newly-cicatrized parts, they had not been sufficient to resist the strong tendency to relapse which characterizes these cases. A certain amount of contraction had again taken place, and two folds of skin, occupying the place of the two natural folds of the axilla, stood out in relief from the surface, and prevented complete extension at the shoulder-joint and also at the elbow. In the hope of remedying this defect, and obtaining a more per-

fect result, a second operation was performed on the 1st of December.

Second Operation.—The anterior axillary fold of skin being gathered up between the thumb and fingers, and drawn out from the body, its base was transfixed at its lower part with a sharp-pointed, straight bistoury, and an incision carried upward toward the axilla and outward through the surface. A tongue-shaped patch of skin was thus detached, which, left to itself, immediately receded and spread itself out upon the chest, thus permitting the arm again to be elevated to a vertical position. In order to close the newly-made wound, and relieve the sutures that were to unite its opposite edges, an incision four inches long was made parallel with, and three fingers' breadths distant from, the wound on either side of it. The edges of these incisions gaped widely apart and afforded the desired relief, and the raw surfaces thus exposed were left to heal by granulation after being covered by a collodion-crust. Complete cicatrization of all the parts involved in the second operation was obtained, under the same management as was employed after the first operation. The elevated position of the arm and suitable gymnastic exercises were also perseveringly enforced.

Though a further improvement was obtained by this second operation, there remained still, in the situation of the anterior axillary fold, an elevated

fold of skin which was prolonged downward upon the arm, extending over the elbow, and also down on the forearm in the shape of a tense-corded band. When complete extension of the arm was attempted, resistance was made by this corded band. Believing that this remaining resistance could be overcome, I determined on a third operation, which was performed March 15, 1875, as follows :

Third Operation.—The fold of skin in the axilla being gathered up between the thumb and fingers, was divided across at two points two inches apart, and the incisions extended through the sound skin a short distance on either side. Between the elbow and shoulder the cicatricial band was also divided across at two points. After relaxation thus obtained, the extension of the limb could be carried to the utmost degree without any remaining resistance. The same treatment of the newly-made wound-surfaces and the same management of the limb itself were continued as after the previous operations.

April 28th.—Cicatrization being again completed with a satisfactory result, an expedient was resorted to which, it was hoped, would permanently resist and eventually overcome entirely the tendency to recontraction on the part of the axillary fold. It consisted of two tubular rings made of stout linen and stuffed with bran, and adapted to be worn high up over the shoulders. These rings, when applied

and drawn toward each other upon the back by means of an elastic strap and buckle, exerted a constant pressure upon the axillary folds in front, and



FIG. 86.

resisted the contraction and consequent elevation of the fold. A trough-like splint, incasing the arm and forearm, and maintaining the elbow in a straight position, was also worn constantly.

The final result of the treatment at the date of his discharge from the hospital (August 17, 1875), was as follows: The limb was capable of the utmost freedom of motion, and, when raised to a vertical position with the elbow and wrist fully extended, there remained no longer any resistance on the part of the fold in the axilla, or along the line of the original cicatricial band that passed down over the bend of the elbow. The shoulder-rings, which are still worn, had effectually prevented any contraction or elevation of the fold in the axilla, and had maintained it in a soft and pliable condition.

Fig. 86, from a photograph taken in July, shows this result.

THE END.

DESCRIPTIVE CATALOGUE

OF

MEDICAL WORKS.



INDEX OF SUBJECTS.

	PAGE		PAGE
Anatomy.....	15	Midwifery.....	25, 26
Anaesthesia.....	25	Mineral Springs.....	29
Acne.....	31	Neuralgia.....	3
Body and Mind.....	17	Nervous System.....	11, 12
Breath, and Diseases which give it a Fetid Odor.....	15	Nursing.....	22
Cerebral Convulsions.....	7	Obstetrics.....	4, 7, 25
Chemical Examination of the Urine in Disease.....	8	Ovarian Tumors.....	23
Chemical Analysis.....	13	" Diagnosis and Treatment.....	30
" Technology.....	30	Paralysis from Brain-Disease.....	31
Chemistry of Common Life.....	16	Physiology.....	9, 10
Clinical Electro-Therapeutics.....	10	Physiology of Common Life.....	16
" Lectures and Essays.....	31	Physiology and Pathology of the Mind.....	17
Comparative Anatomy.....	6	Physiological Effects of Severe Muscular Exercise.....	11
Club-foot.....	24	Pulmonary Consumption.....	5
Diseases of the Nervous System.....	11	Practical Medicine.....	20
" " Bones.....	18	Physical Cause of the Death of Christ.....	25
" " Women.....	25, 26	Popular Science.....	32
" " the Chest.....	25	Puerperal Diseases.....	2
" " Children.....	24, 28	Reports.....	4
" " the Rectum.....	23	Recollections of Past Life.....	14
" " the Ovaries.....	30	" of the Army of the Potomac.....	16
Emergencies.....	14	Responsibility in Mental Diseases.....	18
Electricity and Practical Medicine.....	19	Sea-Sickness.....	2
Foods.....	24	Surgical Pathology.....	5
Galvano-Therapeutics.....	22	" Diseases of the Male Genito-Urinary Organs.....	27
Hospitalism.....	25	Surgery.....	7
Histology and Histo-Chemistry of Man.....	8	" Orthopedic.....	31
Infancy.....	6	Syphilis.....	27
Insanity in its Relation to Crime.....	10	Science.....	30, 32
Materia Medica and Therapeutics.....	22	Skin-Diseases.....	21
Medical Journal.....	32	Therapeutics.....	31
Mental Physiology.....	5	Uterine Therapeutics.....	26
		Winter and Spring.....	4

D. APPLETON & CO.,
PUBLISHERS AND IMPORTERS,
549 & 551 BROADWAY, NEW YORK.

1876.

CATALOGUE
OF
MEDICAL WORKS.

ANSTIE.

Neuralgia, and Diseases which resemble it.

By FRANCIS E. ANSTIE, M. D., F. R. C. P.,

Senior Assistant Physician to Westminster Hospital; Lecturer on Materia Medica in Westminster Hospital School; and Physician to the Belgrave Hospital for Children; Editor of "The Practitioner" (London), etc.

1 vol., 12mo. Cloth, \$2.50.

"It is a valuable contribution to scientific medicine."—*The Lancet* (London).

BARKER.

The Puerperal Diseases. *Clinical Lectures
delivered at Bellevue Hospital.*

By FORDYCE BARKER, M. D.,

Clinical Professor of Midwifery and the Diseases of Women in the Bellevue Hospital Medical College; Obstetric Physician to Bellevue Hospital; Consulting Physician to the New York State Woman's Hospital; Fellow of the New York Academy of Medicine; formerly President of the Medical Society of the State of New York; Honorary Fellow of the Obstetrical Societies of London and Edinburgh; Honorary Fellow of the Royal Medical Society of Athens, Greece, etc., etc., etc.

Third Edition. 1 vol., 8vo. Cloth. 526 pages. Price, \$5; Sheep, \$6.

"For nearly twenty years it has been my duty, as well as my privilege, to give clinical lectures at Bellevue Hospital, on midwifery, the puerperal and the other diseases of women. This volume is made up substantially from phonographic reports of the lectures which I have given on the puerperal diseases. Having had rather exceptional opportunities for the study of these diseases, I have felt it to be an imperative duty to utilize, so far as lay in my power, the advantages which I have enjoyed for the promotion of science, and, I hope, for the interests of humanity. In many subjects, such as albuminuria, convulsions, thrombosis, and embolism, septicæmia, and pyæmia, the advance of science has been so rapid as to make it necessary to teach something new every year. Those, therefore, who have formerly listened to my lectures on these subjects, and who now do me the honor to read this volume, will not be surprised to find, in many particulars, changes in pathological views, and often in therapeutical teaching, from doctrines before inculcated. At the present day, for the first time in the history of the world, the obstetric department seems to be assuming its proper position, as the highest branch of medicine, if its rank be graded by its importance to society, or by the intellectual culture and ability required, as compared with that demanded of the physician or the surgeon. A man may become eminent as a physician, and yet know very little of obstetrics; or he may be a successful and distinguished surgeon, and be quite ignorant of even the rudiments of obstetrics. But no one can be a really able obstetrician unless he be both physician and surgeon. And, as the greater includes the less, obstetrics should rank as the highest department of our profession."—*From Author's Preface.*

On Sea-sickness.

By FORDYCE BARKER, M. D.

1 vol., 16mo. 36 pp. Flexible Cloth, 75 cents.

Reprinted from the NEW YORK MEDICAL JOURNAL. By reason of the great demand for the number of that journal containing the paper, it is now presented in book form, with such prescriptions added as the author has found useful in relieving the suffering from sea-sickness.

BARNES.

Obstetric Operations, *including the Treatment of Hæmorrhage, and forming a Guide to the Management of Difficult Labor.*

By ROBERT BARNES, M. D., London, F. R. C. P.,

Obstetric Physician and Lecturer on Obstetrics and the Diseases of Women and Children to St. George's Hospital; Examiner in Obstetrics to the Royal College of Physicians and the Royal College of Surgeons; President of the Metropolitan Branch of the British Medical Association; late Examiner to the University of London; formerly Obstetric Physician to the London and to St. Thomas Hospitals; and late Physician to the Eastern Division of the Royal Maternity Charity.

Third Edition. Revised and extended. 1 vol., 8vo. 606 pages. Cloth, \$4.50.

"Such a work as Dr. Barnes's was greatly needed. It is calculated to elevate the practice of the obstetric art in this country, and to be of great service to the practitioner."—*Lancet*.

"The book of Dr. Barnes is not, properly speaking, a dogmatic treatise on obstetric operations. It is a series of original lectures, comprising, at one and the same time, a practical analysis of the serious accidents in parturition, the reasoned-out indications for, and the most judicious researches in the manner of operating, the method to choose, the instrument to prefer, and the details of the manœuvres required to insure success. The clearness of the style is perfect. The order, without being altogether rigorous, is what it is able to be generally in a series of clinical lectures. The description of the instruments, the application of the forceps, cephalotripsy, embryotomy, Cæsarean section, the practical reflections on narrowing and malformation of the pelvis, ruptures of the uterus, placenta prævia, hæmorrhage, and, in fact, all the grand questions in obstetrics are treated with accurate good sense. At each instant, by some remark or other, is revealed a superior mind, ripened by having seen much and meditated much."—*From Preface to the French Edition by Prof. Pajot.*

BASTIAN.

Paralysis from Brain Disease in its *Common Forms.*

By H. CHARLTON BASTIAN, M. A., M. D., F. R. S.,

Fellow of the Royal College of Physicians; Professor of Pathological Anatomy in University College, London; Physician to University College Hospital; and Senior Assistant Physician to the National Hospital for the Paralyzed and Epileptic.

With Illustrations. 1 vol., 12mo. Cloth. \$1.75.

PREFACE.

These Lectures were delivered in University College Hospital last year, at a time when I was doing duty for one of the senior physicians, and during the same year—after they had been reproduced from very full notes taken by my friend Mr. John Tweedy—they appeared in the pages of *The Lancet*.

They are now republished at the request of many friends, though only after having undergone a very careful revision, during which a considerable quantity of new matter has been added. It would have been easy to have very much increased the size of the book by the introduction of a larger number of illustrative cases, and by treatment of many of the subjects at greater length, but this the author has purposely abstained from doing, under the belief that in its present form it is likely to prove more acceptable to students, and also perhaps more useful to busy practitioners.

Notwithstanding its defects and many shortcomings, the author is not without a hope that this little book may be considered in some measure to supply a deficiency which has long existed in medical literature. No department of medicine stands more in need of being represented in a text book of moderate compass; so that, imperfect as it is, this small work may perhaps be of some service till it is superseded by something better. In it the author has endeavored to treat the subject with more precision than has hitherto been customary, and, while the lectures contain some novelties in method and mode of exposition, he hopes they may also be found not unfaithfully to embody the principal facts at present known concerning this very important class of diseases.

BENNET.

On the Treatment of Pulmonary Consumption, by Hygiene, Climate, and Medicine, in its Connection with Modern Doctrines.

BY JAMES HENRY BENNET, M. D.,

Member of the Royal College of Physicians, London; Doctor of Medicine of the University of Paris, etc., etc.

1 vol., thin 8vo. Cloth, \$1.50.

An interesting and instructive work, written in the strong, clear, and lucid manner which appears in all the contributions of Dr. Bennet to medical or general literature.

"We cordially commend this book to the attention of all, for its practical common-sense views of the nature and treatment of the scourge of all temperate climates, pulmonary consumption."—*Detroit Review of Medicine*.

Winter and Spring on the Shores of the Mediterranean; or, the Riviera, Mentone, Italy, Corsica, Sicily, Algeria, Spain, and Biarritz, as Winter Climates.

This work embodies the experience of ten winters and springs passed by Dr. Bennet on the shores of the Mediterranean, and contains much valuable information for physicians in relation to the health-restoring climate of the regions described.

1 vol., 12mo. 621 pp. Cloth, \$3.50.

"Exceedingly readable, apart from its special purposes, and well illustrated."—*Evening Commercial*.

"It has a more substantial value for the physician, perhaps, than for any other class or profession. . . . We commend this book to our readers as a volume presenting two capital qualifications—it is at once entertaining and instructive."—*N. Y. Medical Journal*.

BILLROTH.

General Surgical Pathology and Therapeutics, in Fifty Lectures. A Text-book for Students and Physicians.

BY DR. THEODOR BILLROTH.

Translated from the Fifth German Edition, with the special permission of the Author, by

CHARLES E. HACKLEY, A. M., M. D.,

Surgeon to the New York Eye and Ear Infirmary; Physician to the New York Hospital; Fellow of the New York Academy of Medicine, etc.

1 vol., 8vo. 714 pp., and 152 Woodcuts. Cloth, \$5.00; Sheep, \$3.00.

Professor Theodor Billroth, one of the most noted authorities on Surgical Pathology, gives in this volume a complete résumé of the existing state of knowledge in this branch of medical science. The fact of this publication going through four editions in Germany, and having been translated into French, Italian, Russian, and Hungarian, should be some guarantee for its standing.

"The want of a book in the English language, presenting in a concise form the views of the German pathologists, has long been felt; and we venture to say no book could more perfectly supply that want than the present volume. . . . We would strongly recommend it to all who take any interest in the progress of thought and observation in surgical pathology, and surgery."—*The Lancet*.

"We can assure our readers that they will consider neither money wasted in its purchase, nor time in its perusal."—*The Medical Investigator*.

CARPENTER.

Principles of Mental Physiology, with their Applications to the Training and Discipline of the Mind and the Study of its Morbid Conditions.

BY WM. B. CARPENTER, M. D., LL. D., F. R. S., F. L. S., F. G. S.,

Registrar of the University of London; Corresponding Member of the Institute of France and of the American Philosophical Society, etc.

1 vol., 8vo. Price, \$3.00.

"Among the numerous eminent writers this country has produced, none are more deserving of praise for having attempted to apply the results of Physiological Research to the explanation of the mutual relations of the mind and body than Dr. Carpenter. To him belongs the merit of having scientifically studied and of having in many instances supplied a rational explanation of those phenomena which, under the names of mesmerism, spirit-rapping, electro-biology, and hypnotism, have attracted so large an amount of attention during the last twenty years. . . . We must conclude by recommending Dr. Carpenter's work to the members of our own profession as applying many facts, that have hitherto stood isolated, to the explanation of the functions of the brain and to psychological processes generally."—*The Lancet*.

COMBE.

The Management of Infancy, *Physiological and Moral. Intended chiefly for the Use of Parents.*

BY ANDREW COMBE, M. D.

REVISED AND EDITED

BY SIR JAMES CLARK, K. C. B., M. D., F. R. S.,
Physician-in-ordinary to the Queen.

First American from the Tenth London Edition. 1 vol., 12mo. 302 pp.
Cloth, \$1.50.

"This excellent little book should be in the hand of every mother of a family; and, if some of our lady friends would master its contents, and either bring up their children by the light of its teachings, or communicate the truths it contains to the poor by whom they are surrounded, we are convinced that they would effect infinitely more good than by the distribution of any number of tracts whatever. . . . We consider this work to be one of the few popular medical treatises that any practitioner may recommend to his patients; and, though, if its precepts are followed, he will probably lose a few guineas, he will not begrudge them if he sees his friend's children grow up healthy, active, strong, and both mentally and physically capable."—*The Lancet*.

CHAUVEAU.

The Comparative Anatomy of the
Domesticated Animals.

By A. CHAUVEAU,

PROFESSOR AT THE LYONS VETERINARY SCHOOL.

Second edition, revised and enlarged, with the coöperation of S. ARLOING, late Principal of Anatomy at the Lyons Veterinary School; Professor at the Toulouse Veterinary School. Translated and edited by GEORGE FLEMING, F. R. G. S., M. A. I., Veterinary Surgeon, Royal Engineers.

1 vol., 8vo. Cloth. 957 pp., with 450 illustrations. Price, \$6.00.

OPINIONS OF THE PRESS.

"This is a valuable work, well conceived and well executed by the authors, MM. Chauveau and Arloing, and well translated by Mr. Fleming. It is rather surprising how few works exist, in any language, in which the anatomy of the commoner animals, domestic and otherwise, is given with any approach to detail. Systematic works there are in abundance, but, if the student be desirous of ascertaining any particular point, such as the position and branches of the pneumogastric or sympathetic nerves, or the homologue of a given muscle in several different animals, he may search all day ere he find it. The work before us appears to be well adapted to meet this difficulty.

"The illustrations are very numerous, and Mr. Fleming has introduced a large number that are not contained in the original work.

"Taking it altogether, the book is a very welcome addition to English literature, and great credit is due to Mr. Fleming for the excellence of the translation, and the many additional notes he has appended to Chauveau's treatise."—*Lancet (London)*.

"The want of a text-book on the Comparative Anatomy of the Domesticated Animals has long been felt. . . . The descriptions of the text are illustrated and assisted by no less than 450 excellent woodcuts. In a work which ranges over so vast a field of anatomical detail and description, it is difficult to select any one portion for review, but our examination of it enables us to speak in high terms of its general excellence. . . . The care and attention with which hipponomy has been cultivated on the Continent are illustrated by every page in M. Chauveau's work."—*Medical Times and Gazette (London)*.

DAVIS.

**Conservative Surgery, as exhibited in remedying
some of the Mechanical Causes that operate injuriously
both in Health and Disease. With Illustrations.**

By HENRY G. DAVIS, M. D.,

Member of the American Medical Association, etc., etc.

1 vol., 8vo. 315 pp. Cloth, \$3.00.

The author has enjoyed rare facilities for the study and treatment of certain classes of disease, and the records here presented to the profession are the gradual accumulation of over thirty years' investigation.

"Dr. Davis, bringing, as he does to his specialty, a great aptitude for the solution of mechanical problems, takes a high rank as an orthopedic surgeon, and his very practical contribution to the literature of the subject is both valuable and opportune. We deem it worthy of a place in every physician's library. The style is unpretending, but trenchant, graphic, and, best of all, quite intelligible."—*Medical Record*.

ECKER.

**The Cerebral Convulsions of Man,
represented according to Personal Investigations, espe-
cially on their Development in the Fœtus, and with ref-
erence to the Use of Physicians.**

By ALEXANDER ECKER,

Professor of Anatomy and Comparative Anatomy in the University of Freiburg.

Translated from the German by Robert T. Edes, M. D.

1 vol., 8vo. 87 pp. \$1.25.

"The work of Prof. Ecker is noticeable principally for its succinctness and clearness, avoiding long discussions on undecided points, and yet sufficiently furnished with references to make easy its comparisons with the labors of others in the same direction.

"Entire originality in descriptive anatomy is out of the question, but the facts verified by our author are here presented in a more intelligible manner than in any other easily-accessible work.

"The knowledge to be derived from this work is not furnished by any other text-book in the English language."—*Boston Medical and Surgical Journal*, January 20, 1873.

ELLIOT.

**Obstetric Clinic. A Practical Contribution to the Study
of Obstetrics, and the Diseases of Women and Children.**

By the late GEORGE T. ELLIOT, M. D.,

Late Professor of Obstetrics and Diseases of Women and Children in the Bellevue Hospital Medical College; Physician to Bellevue Hospital and to the New York Lying-in Asylum; Consulting Physician to the Nursery and Child's Hospital; Consulting Surgeon to the State Woman's Hospital; Corresponding Member of the Edinburgh Obstetrical Society and of the Royal Academy of Havana; Fellow of the N. Y. Academy of Medicine; Member of the County Medical Society, of the Pathological Society, etc., etc.

1 vol., 8vo. 458 pp. Cloth, \$4.50.

This work is, in a measure, a *résumé* of separate papers previously prepared by the late Dr. Elliot; and contains, besides, a record of nearly two hundred important and difficult cases in midwifery, selected from his own practice. It has met with a hearty reception, and has received the highest encomiums both in this country and in Europe.

FREY.

The Histology and Histo-Chemistry
of Man. A Practical Treatise on the Elements of Composition and Structure of the Human Body.

By HEINRICH FREY,
Professor of Medicine in Zurich.

Translated from the Fourth German Edition, by Arthur E. J. Barker,
Surgeon to the City of Dublin Hospital; Demonstrator of Anatomy, Royal College of Surgeons,
Ireland; Visiting Surgeon, Convalescent Home, Stillorgan; and revised by the Author.
With 680 Engravings.

1 vol., 8vo. Cloth, \$5.00; Sheep, \$6.00.

CONTENTS.

The Elements of Composition and of Structure of the Body: Elements of Composition—Albuminous or Protein Compounds, Hæmoglobin, Histogenic Derivatives of the Albuminous Substances or Albuminoids, the Fatty Acids and Fats, the Carbohydrates, Non-Nitrogenous Acids, Nitrogenous Acids, Amides, Amido-Acids, and Organic Bases, Animal Coloring Matters, Cyanogen Compounds, Mineral Constituents; Elements of Structure—the Cell, the Origin of the Remaining Elements of Tissue; the Tissues of the Body—Tissues composed of Simple Cells, with Fluid Intermediate Substance, Tissues composed of Simple Cells, with a small amount of Solid Intermediate Substance, Tissues belonging to the Connective-Substance Group, Tissues composed of Transformed, and, as a rule, Cohering Cells, with Homogeneous, Scanty, and more or less Solid Intermediate Substance, Composite Tissues: The Organs of the Body—Organs of the Vegetative Type, Organs of the Animal Group.

FLINT.

Manual of Chemical Examination of
the Urine in Disease. With Brief Directions for the Examination of the most Common Varieties of Urinary Calculi.

By AUSTIN FLINT, JR., M. D.,

Professor of Physiology and Microscopy in the Bellevue Hospital Medical College; Fellow of the New York Academy of Medicine; Member of the Medical Society of the County of New York; Resident Member of the Lyceum of Natural History in the City of New York, etc.

Third Edition, revised and corrected. 1 vol., 12mo. 77 pp. Cloth, \$1.00.

The chief aim of this little work is to enable the busy practitioner to make for himself, rapidly and easily, all ordinary examinations of Urine; to give him the benefit of the author's experience in eliminating little difficulties in the manipulations, and in reducing processes of analysis to the utmost simplicity that is consistent with accuracy.

"We do not know of any work in English so complete and handy as the Manual now offered to the Profession by Dr. Flint, and the high scientific reputation of the author is a sufficient guarantee of the accuracy of all the directions given."—*Journal of Applied Chemistry.*

"We can unhesitatingly recommend this Manual."—*Psychological Journal.*

"Eminently practical."—*Detroit Review of Medicine.*

FLINT.

The Physiology of Man. *Designed to represent the Existing State of Physiological Science as applied to the Functions of the Human Body.*

By AUSTIN FLINT, JR., M. D.,

Professor of Physiology and Microscopy in the Bellevue Hospital Medical College, and in the Long Island College Hospital; Fellow of the New York Academy of Medicine; Microscopist to Bellevue Hospital.

New and thoroughly revised Edition. In Five Volumes. 8vo. Tinted Paper.

Volume I.—*The Blood; Circulation; Respiration.*

8vo. 502 pp. Cloth, \$4.50.

"If the remaining portions of this work are compiled with the same care and accuracy, the whole may vie with any of those that have of late years been produced in our own or in foreign languages."—*British and Foreign Medico-Chirurgical Review.*

"As a book of general information it will be found useful to the practitioner, and, as a book of reference, invaluable in the hands of the anatomist and physiologist."—*Dublin Quarterly Journal of Medical Science.*

"The complete work will prove a valuable addition to our systematic treatises on human physiology."—*The Lancet.*

"To those who desire to get in one volume a concise and clear, and at the same time sufficiently full *résumé* of 'the existing state of physiological science,' we can heartily recommend Dr. Flint's work. Moreover, as a work of typographical art it deserves a prominent place upon our library-shelves. Messrs. Appleton & Co. deserve the thanks of the profession for the very handsome style in which they issue medical works. They give us hope of a time when it will be very generally believed by publishers that physicians' eyes are worth saving."—*Medical Gazette.*

Volume II.—*Alimentation; Digestion; Absorption; Lymph and Chyle.*

8vo. 556 pp. Cloth, \$4.50.

"The second instalment of this work fulfils all the expectations raised by the perusal of the first. . . . The author's explanations and deductions bear evidence of much careful reflection and study. . . . The entire work is one of rare interest. The author's style is as clear and concise as his method is studious, careful, and elaborate."—*Philadelphia Inquirer.*

"We regard the two treatises already issued as the very best on human physiology which the English or any other language affords, and we recommend them with thorough confidence to students, practitioners, and laymen, as models of literary and scientific ability."—*N. Y. Medical Journal.*

"We have found the style easy, lucid, and at the same time terse. The practical and positive results of physiological investigation are succinctly stated, without, it would seem, extended discussion of disputed points."—*Boston Medical and Surgical Journal.*

"It is a volume which will be welcome to the advanced student, and as a work of reference."—*The Lancet.*

"The leading subjects treated of are presented in distinct parts, each of which is designed to be an exhaustive essay on that to which it refers."—*Western Journal of Medicine.*

Flint's Physiology. Volume III.—*Secretion; Excretion; Ductless Glands; Nutrition; Animal Heat; Movements; Voice and Speech.*

8vo. 526 pp. Cloth, \$4.50.

"Dr. Flint's reputation is sufficient to give a character to the book among the profession, where it will chiefly circulate, and many of the facts given have been verified by the author in his laboratory and in public demonstration."—*Chicago Courier.*

"The author bestows judicious care and labor. Facts are selected with discrimination, theories critically examined, and conclusions enunciated with commendable clearness and precision."—*American Journal of the Medical Sciences.*

Volume IV.—*The Nervous System.*

8vo. Cloth, \$4.50.

This volume embodies the results of exhaustive study, and of a long and laborious series of experiments, presented in a manner remarkable for its strength and clearness. No other department of physiology has so profound an interest for the modern and progressive physician as that pertaining to the nervous system. The diseases of this system are now engaging the study and attention of some of the greatest minds in the medical world, and in order to follow their brilliant discoveries and developments, especially in connection with the science of electrology, it is absolutely necessary to obtain a clear and settled knowledge of the anatomy and physiology of the nervous system. It is the design of this work to impart that knowledge free from the perplexing speculations and uncertainties that have no real value for the practical student of medicine. The author boldly tests every theory for himself, and asks his readers to accept nothing that is not capable of demonstration. The properties of the cerebro-spinal, nervous, and sympathetic systems are treated of in a manner at once lucid, thorough, and interesting.

Although this volume is one, perhaps the most important one, of the author's admirable series in the *Physiology of Man*, it is nevertheless complete in itself, and may be safely pronounced indispensable to every physician who takes a pride and interest in the progress of medical science.

Volume V.—*Special Senses; Generation.*

8vo. Cloth, \$4.50.

"The present volume completes the task, begun eleven years ago, of preparing a work, intended to represent the existing state of physiological science, as applied to the functions of the human body. The kindly reception which the first four volumes have received has done much to sustain the author in an undertaking, the magnitude of which he has appreciated more and more as the work has progressed.

"In the fifth and last volume, an attempt has been made to give a clear account of the physiology of the special senses and generation, a most difficult and delicate undertaking. . . .

"Finally, as regards the last, as well as the former volumes, the author can only say that he has spared neither time nor labor in their preparation; and the imperfections in their execution have been due to deficiency in ability and opportunity. He indulges the hope, however, that he has written a book which may assist his fellow-workers, and interest, not only the student and practitioner of medicine, but some others who desire to keep pace with the progress of Natural Science."—*Extracts from Preface.*

Flint's Text-Book of Human Physiology, for the Use of Students and Practitioners of Medicine.

In one large octavo volume of 978 pages, elegantly printed on fine paper, and profusely illustrated with three Lithographic Plates and 313 Engravings on Wood. Price, in cloth, \$6.00; sheep, \$7.00.

While Prof. Flint's "Physiology of Man," in five octavo volumes, also published by D. Appleton & Co., is invaluable as a book of reference, giving an epitome of the literature of physiology, with copious references to other authors, the publishers have appreciated the necessity for a new text-book, for the use of students and practitioners of medicine.

This new work is intended to meet this pressing want, and it contains most of the facts presented in the larger treatise, without historical references or discussions of minor and controverted questions. The high reputation of the author as a public teacher, and the success of the larger treatise, render it certain that the "Text-book" will be admirably adapted to the wants of medical students.

In the "Text-book," all important points connected with Human Physiology are treated of fully and clearly, and many subjects, such as the Nervous System, the Special Senses, etc., the treatment of which is barren and unsatisfactory in many works written or republished in this country, are brought fully up to the requirements of the day.

The publishers have given great attention to the execution of the illustrations, few of which are familiar to American readers. It being almost impossible to reproduce some of the cuts taken from foreign works, they have succeeded in obtaining abroad about one hundred electrotypes from the original engravings contained in Sappey's great work upon Anatomy, which are unequaled in their mechanical execution. The subject of Generation is also illustrated by lithographic plates taken from Haeckel.

The great care necessary in the printing of the elaborate illustrations has caused an unavoidable delay in the appearance of the work; but the publishers feel confident that it will fully meet their expectations, and justify the reputation of its author.

"In preparing this text-book for the use of students and practitioners of medicine, I have endeavored to adapt it to the wants of the profession, as they have appeared to me after a considerable experience as a public teacher of human physiology. My large treatise in five volumes is here condensed, and I have omitted bibliographical citations and matters of purely historical interest. Many subjects, which were considered rather elaborately in my larger work, are here presented in a much more concise form. I have added, also, numerous illustrations, which I hope may lighten the labors of the student. A few of these are original, but by far the greatest part has been selected from reliable authorities. I have thought it not without historical interest to reproduce exactly some of the classical engravings from the works of great discoverers, such as illustrations contained in the original editions of Fabricius, Harvey, and Asellius. In addition, I have reproduced a few of the beautiful microscopical photographs taken at the United States Army Medical Museum, under the direction of Dr. J. J. Woodward, to whom I here express my grateful acknowledgments. I have also to thank M. Sappey for his kindness in furnishing electrotypes of many of the superb engravings with which his great work upon Anatomy is illustrated.

"My work in five volumes was intended as a book of reference, which I hope will continue to be useful to those who desire an account of the literature of physiology, as well as a statement of the facts of the science. I have always endeavored, in public teaching, to avoid giving undue prominence to points in which I might myself be particularly interested, from having made them subjects of special study or of original research. In my text-book I have carried out the same idea, striving to teach, systematically and with uniform emphasis, what students of medicine are expected to learn in physiology, and avoiding elaborate discussions of subjects not directly connected with practical medicine, surgery, and obstetrics. While I have referred to my original observations upon the location of the sense of want of air in the general system, the new excretory function of the liver, the function of glycogenesis, the influence of muscular exercise upon the elimination of uric acid, etc., I have not considered these subjects with great minuteness, and have generally referred the reader to monographs for the details of my experiments.

"Finally, in presenting this work to the medical profession, I cannot refrain from an expression of my acknowledgments to the publishers, who have spared nothing in carrying out my views, and have devoted special pains to the mechanical execution of the illustrations."—*Author's Preface.*

FLINT.

On the Physiological Effects of Severe and Protracted Muscular Exercise. *With special refer- ence to its Influence upon the Excretion of Nitrogen.*

By AUSTIN FLINT, JR., M. D.,

Professor of Physiology in the Bellevue Hospital Medical College, New York, etc., etc.

1 vol., 8vo. 91 pp. Cloth, \$2.00.

This monograph on the relations of Urea to Exercise is the result of a thorough and careful investigation made in the case of Mr. Edward Payson Weston, the celebrated pedestrian. The chemical analyses were made under the direction of R. O. Doremus, M. D., Professor of Chemistry and Toxicology in the Bellevue Hospital Medical College, by Mr. Oscar Loew, his assistant. The observations were made with the coöperation of J. C. Dalton, M. D., Professor of Physiology in the College of Physicians and Surgeons; Alexander B. Mott, M. D., Professor of Surgical Anatomy; W. H. Van Buren, M. D., Professor of Principles of Surgery; Austin Flint, M. D., Professor of the Principles and Practice of Medicine; W. A. Hammond, M. D., Professor of Diseases of the Mind and Nervous System—all of the Bellevue Hospital Medical College.

"This work will be found interesting to every physician. A number of important results were obtained valuable to the physiologist."—*Cincinnati Medical Repertory.*

HAMILTON.

Clinical Electro-Therapeutics. (*Medical and Surgical.*) *A Manual for Physicians for the Treatment more especially of Nervous Diseases.*

By ALLAN McLANE HAMILTON, M. D.,

Physician in charge of the New York State Hospital for Diseases of the Nervous System; Member of the New York Neurological and County Medical Societies, etc., etc.

With Numerous Illustrations. 1 vol., 8vo. Cloth. Price, \$2.00.

This work is the compilation of well-tried measures and reported cases, and is intended as a simple guide for the general practitioner. It is as free from confusing theories, technical terms, and unproved statements, as possible. Electricity is indorsed as a very valuable remedy in certain diseases, and as an invaluable therapeutical means in nearly all forms of NERVOUS DISEASE; but not as a specific for every human ill, mental and physical.

HAMMOND.

Insanity in its Relations to Crime. *A Text and a Commentary.*

By WILLIAM A. HAMMOND, M. D.

1 vol., 8vo. 77 pp. Cloth, \$1.00.

"A part of this essay, under the title 'Society versus Insanity,' was contributed to *Putnam's Magazine*, for September, 1870. The greater portion is now first published. The importance of the subject considered can scarcely be over-estimated, whether we regard it from the stand-point of science or social economy; and, if I have aided in its elucidation, my object will have been attained."—*From Author's Preface.*

Clinical Lectures on Diseases of the Nervous System. *Delivered at the Bellevue Hospital Medical College.*

By WILLIAM A. HAMMOND, M. D.,

Professor of Diseases of the Mind and Nervous System, etc. Edited, with Notes, by T. M. B. CROSS, M. D., Assistant to the Chairs of Diseases of the Mind and Nervous System, etc.

In one handsome volume of 300 pages. Price, \$3.50.

HAMMOND.

A Treatise on Diseases of the Nervous
System.

BY WILLIAM A. HAMMOND, M. D.,

Professor of Diseases of the Mind and Nervous System in the Medical Department of the University of the City of New York; President of the New York Neurological Society, etc., etc.

Sixth Edition. 1 vol., 8vo. Strong Cloth Binding, \$6.00; Sheep, \$7.00.

The remarkable success attendant on the issue of the five previous editions of this work in less than four years has encouraged the author and publishers to attempt to make the work still more worthy the confidence of the medical profession. A great part of the treatise has been entirely rewritten, and several new chapters have been added. By a change in type, and enlarging the page, the new matter, amounting to one-half of the original work, has been added without increasing materially the bulk of the volume. Many new illustrations have been incorporated in the text, and the whole treatise has been brought fully up to the present time. In addition to the fund of personal observation and experience adduced by Prof. Hammond, the labors of English, French, and German writers have received due attention.

Among the diseases considered in the present edition, which were not treated of in the former editions, are: Chronic Verticalar Meningitis; Chronic Basilar Meningitis; Cervical Pachy-Meningitis; Spinal Paralysis of Adults; Amyotrophic Lateral Spinal Sclerosis; Facial Atrophy; Organic Diseases of Nerves; Chronic Alcoholic Intoxication; Delirium Tremens; Exophthalmic Goitre; and Anapeiratic Paralysis—paralysis induced by a frequent repetition of certain muscular actions. Besides which, extensive alterations and additions have been made to the remarks on other affections—the departments of Morbid Anatomy, Pathology, and Treatment, being especially amplified.

NOTICES OF FORMER EDITIONS.

"Free from useless verbiage and obscurity, it is evidently the work of a man who knows what he is writing about, and knows how to write about it."—*Chicago Medical Journal.*

"Unquestionably the most complete treatise on the diseases to which it is devoted which has yet appeared in the English language."—*London Medical Times and Gazette.*

"This is a valuable and comprehensive book; it embraces many topics, and extends over a wide sphere. One of the most valuable parts of it relates to the Diseases of the Brain; while the remaining portion of the volume treats of the Diseases of the Spinal Cord, the Cerebro-spinal System, the Nerve-Cells, and the Peripheral Nerves."—*British Medical Journal.*

"The work before us is unquestionably the most exhaustive treatise, on the diseases to which it is devoted, that has yet appeared in English. And its distinctive value arises from the fact that the work is no mere *rafficiamento* of old observations, but rests on his own experience and practice, which, as we have before observed, have been very extensive."—*American Journal of Syphilography.*

"The author of this work has attained a high rank among our brethren across the Atlantic from previous labors in connection with the disorders of the nervous system, as well as from various other contributions to medical literature, and he now holds the official appointments of Physician to the New York State Hospital for the Diseases of the Nervous System, and Professor of the same department in the Bellevue Hospital Medical College. The present treatise is the fruit of the experience thus acquired, and we have no hesitation in pronouncing it a most valuable addition to our systematic literature."—*Glasgow Medical Journal.*


HOFFMANN.

Manual of Chemical Analysis, as applied to the Examination of Medicinal Chemicals and their Preparations. A Guide for the Determination of their Identity and Quality, and for the Detection of Impurities and Adulterations. For the use of Pharmacutists, Physicians, Druggists, and Manufacturing Chemists, and Pharmaceutical and Medical Students.

By FRED. HOFFMANN, PHIL. D.

One vol., 8vo. Richly Illustrated. Cloth. Price, \$3.

SPECIMEN OF ILLUSTRATIONS.



"This volume is a carefully-prepared work, and well on to the existing state of both the science and art of modern pharmacy. It is a book which will find its place in every medical and pharmaceutical laboratory and library, and is a safe and instructive guide to medical students and practitioners of medicine."—*American Journal of Science and Arts.*

In America this work has already met with general and unqualified approval; and in Europe is now being welcomed as one of the best and most important additions to modern pharmaceutical literature.

Send for descriptive circular. Address

D. APPLETON & CO., 549 & 551 Broadway, N. Y. City.

HOLLAND.

Recollections of Past Life,

By SIR HENRY HOLLAND, Bart., M. D., F. R. S., K. C. B., etc.,
President of the Royal Institution of Great Britain, Physician-in-Ordinary to the Queen,
etc., etc.

1 vol., 12mo, 351 pp. Price, Cloth, \$2.00.

A very entertaining and instructive narrative, partaking somewhat of the nature of autobiography and yet distinct from it, in this, that its chief object, as alleged by the writer, is not so much to recount the events of his own life, as to perform the office of chronicler for others with whom he came in contact and was long associated.

The "Life of Sir Henry Holland" is one to be recollected, and he has not erred in giving an outline of it to the public.—*The Lancet*.

"His memory was—is, we may say, for he is still alive and in possession of all his faculties—stored with recollections of the most eminent men and women of this century. . . . A life extending over a period of eighty-four years, and passed in the most active manner, in the midst of the best society, which the world has to offer, must necessarily be full of singular interest; and Sir Henry Holland has fortunately not waited until his memory lost its freshness before recalling some of the incidents in it."—*The New York Times*.

HOWE.

Emergencies, and How to Treat Them.

The Etiology, Pathology, and Treatment of Accidents, Diseases, and Cases of Poisoning, which demand Prompt Attention. Designed for Students and Practitioners of Medicine.

By JOSEPH W. HOWE, M. D.,

Clinical Professor of Surgery in the Medical Department of the University of New York
Visiting Surgeon to Charity Hospital; Fellow of the New York Academy
of Medicine, etc., etc.

1 vol., 8vo. Cloth, \$3.00.

"This work has a taking title, and was written by a gentlemen of acknowledged ability, to fill a void in the profession. . . . To the general practitioner in towns, villages, and in the country, where the aid and moral support of a consultation cannot be availed of, this volume will be recognized as a valuable help. We commend it to the profession.—*Cincinnati Lancet and Observer*.

"This work is certainly novel in character, and its usefulness and acceptability are as marked as its novelty. . . . The book is confidently recommended."—*Richmond and Louisville Medical Journal*.

"This volume is a practical illustration of the positive side of the physician's life, a constant reminder of what he is to do in the sudden emergencies which frequently occur in practice. . . . The author wastes no words, but devotes himself to the description of each disease as if the patient were under his hands. Because it is a good book we recommend it most heartily to the profession."—*Boston Medical and Surgical Journal*.

"This work bears evidence of a thorough practical acquaintance with the different branches of the profession. The author seems to possess a peculiar aptitude for imparting instruction as well as for simplifying tedious details. . . . A careful perusal will amply repay the student and practitioner."—*New York Medical Journal*.

"This is the best work of the kind we have ever seen."—*New York Journal of Psychological Medicine*.

HOWE.

The Breath, and the Diseases which give
it a Fetid Odor. With Directions for Treatment.

By JOSEPH W. HOWE, M. D.,

author of "Emergencies," "Winter Homes," etc; Clinical Professor of Surgery in the Medical Department of the University of New York; Visiting Surgeon to Charity and St. Francis Hospitals; Fellow of the New York Academy of Medicine, etc.

"It is somewhat remarkable that the subject of fetid breath, which occasions so much annoyance. . . . should have attracted so little attention from authors and investigators. Hence a thoroughly scientific exposition of the whole subject, such as Dr. Howe has given us, has long been a desideratum. . . . This little volume well deserves the attention of physicians, to whom we commend it most highly."—*Chicago Medical Journal*.

". . . To any one suffering from the affection, either in his own person or in that of his intimate acquaintances, we can commend this volume as containing all that is known concerning the subject, set forth in a pleasant style."—*Philadelphia Medical Times*.

"This little work is on a subject that has heretofore been almost entirely ignored by medical authors, yet its importance is well known by every practitioner. . . . The author gives a succinct account of the diseased conditions in which a fetid breath is an important symptom, with his method of treatment. We consider the work a real addition to medical literature."—*Cincinnati Medical Journal*.

HUXLEY AND YOUMANS.

The Elements of Physiology and Hygiene. With Numerous Illustrations.

By THOMAS H. HUXLEY, LL. D., F. R. S., and
WILLIAM JAY YOUMANS, M. D.

New and Revised Edition. 1 vol., 12mo. 420 pp. \$1.75.

A text-book for educational institutions, and a valuable elementary work for students of medicine. The greater portion is from the pen of Professor Huxley, adapted by Dr. Youmans to the circumstances and requirements of American education. The eminent claim of Professor Huxley's "Elementary Physiology" is, that, while up to the times, it is trustworthy in its presentation of the subject; while rejecting discredited doctrines and doubtful speculations, it embodies the latest results that are established, and represents the present actual state of physiological knowledge.

"A valuable contribution to anatomical and physiological science."—*Religious Telescope*.

"A clear and well arranged work, embracing the latest discoveries and accepted theories."—*Buffalo Commercial*.

"Teeming with information concerning the human physical economy."—*Evening Journal*.

HUXLEY.

The Anatomy of Vertebrated Animals.

By THOMAS HENRY HUXLEY, LL. D., F. R. S.,

Author of "Man's Place in Nature," "On the Origin of Species," "Lay Sermons and Addresses," etc.

1 vol., 12mo. Cloth, \$2.50.

The former works of Prof. Huxley leave no room for doubt as to the importance and value of his new volume. It is one which will be very acceptable to all who are interested in the subject of which it treats.

"This long-expected work will be cordially welcomed by all students and teachers of Comparative Anatomy as a compendious, reliable, and, notwithstanding its small dimensions, most comprehensive guide on the subject of which it treats. To praise or to criticise the work of so accomplished a master of his favorite science would be equally out of place. It is enough to say that it realizes, in a remarkable degree, the anticipations which have been formed of it; and that it presents an extraordinary combination of wide, general views, with the clear, accurate, and succinct statement of a prodigious number of individual facts."—*Nature*.

JOHNSON.

The Chemistry of Common Life.

Illustrated with numerous Wood Engravings.

By JAMES F. JOHNSON, M. A., F. R. S., F. G. S., ETC., ETC.,

Author of "Lectures on Agricultural Chemistry and Geology," "A Catechism of Agricultural Chemistry and Geology," etc.

2 vols., 12mo. Cloth, \$3.00.

It has been the object of the author in this work to exhibit the present condition of chemical knowledge, and of matured scientific opinion, upon the subjects to which it is devoted. The reader will not be surprised, therefore, should he find in it some things which differ from what is to be found in other popular works already in his hands or on the shelves of his library.

LETTERMAN.

Medical Recollections of the Army of
the Potomac.

By JONATHAN LETTERMAN, M. D.,

Late Surgeon U. S. A., and Medical Director of the Army of the Potomac.

1 vol., 8vo. 194 pp. Cloth, \$1.00.

"This account of the medical department of the Army of the Potomac has been prepared, amid pressing engagements, in the hope that the labors of the medical officers of that army may be known to an intelligent people, with whom to know is to appreciate; and as an affectionate tribute to many, long my zealous and efficient colleagues, who, in days of trial and danger, which have passed, let us hope never to return, evinced their devotion to their country and to the cause of humanity, without hope of promotion or expectation of reward."—*Preface.*

"We venture to assert that but few who open this volume of medical annals, pregnant as they are with instruction, will care to do otherwise than finish them at a sitting."—*Medical Record.*

"A graceful and affectionate tribute."—*N. Y. Medical Journal.*

LEWES.

The Physiology of Common Life.

By GEORGE HENRY LEWES,

Author of "Seaside Studies," "Life of Goethe," etc.

2 vols., 12mo. Cloth, \$3.00.

The object of this work differs from that of all others on popular science in its attempt to meet the wants of the student, while meeting those of the general reader, who is supposed to be wholly unacquainted with anatomy and physiology.

MAUDSLEY.

The Physiology and Pathology of the
Mind.

BY HENRY MAUDSLEY, M. D., LONDON,

Fellow of the Royal College of Physicians; Professor of Medical Jurisprudence in University College, London; President-elect of the Medico-Psychological Association; Honorary Member of the Medico-Psychological Society of Paris, of the Imperial Society of Physicians of Vienna, and of the Society for the Promotion of Psychiatry and Forensic Psychology of Vienna; formerly Resident Physician of the Manchester Royal Lunatic Asylum, etc., etc.

1 vol., 8vo. 422 pp. Cloth, \$3.00.

This work aims, in the first place, to treat of mental phenomena from a physiological rather than from a metaphysical point of view; and, secondly, to bring the manifold instructive instances presented by the unsound mind to bear upon the interpretation of the obscure problems of mental science.

"Dr. Maudsley has had the courage to undertake, and the skill to execute, what is, at least in English, an original enterprise."—*London Saturday Review*.

"It is so full of sensible reflections and sound truths that their wide dissemination could not but be of benefit to all thinking persons."—*Psychological Journal*.

"Unquestionably one of the ablest and most important works on the subject of which it treats that has ever appeared, and does credit to his philosophical acumen and accurate observation."—*Medical Record*.

"We lay down the book with admiration, and we commend it most earnestly to our readers as a work of extraordinary merit and originality—one of those productions that are evolved only occasionally in the lapse of years, and that serve to mark actual and very decided advantages in knowledge and science."—*N. Y. Medical Journal*.

Body and Mind: *An Inquiry into their Connection and Mutual Influence, especially in reference to Mental Disorders; an enlarged and revised edition to which are added Psychological Essays.*

BY HENRY MAUDSLEY, M. D., LONDON,

Author of "The Physiology and Pathology of the Mind."

1 vol., 12mo. 155 pp. Cloth, \$1.00.

The general plan of this work may be described as being to bring man, both in his physical and mental relations, as much as possible within the scope of scientific inquiry.

"A representative work, which every one must study who desires to know what is doing in the way of real progress, and not mere chatter, about mental physiology and pathology."—*Lancet*.

"It distinctly marks a step in the progress of scientific psychology."—*The Practitioner*.

Responsibility in Mental Diseases.

BY HENRY MAUDSLEY, M. D., LONDON,

Author of "Body and Mind," "Physiology and Pathology of the Mind."

1 vol., 12mo. 313 pp. Cloth, \$1.50.

"This book is a compact presentation of those facts and principles which require to be taken into account in estimating human responsibility—not legal responsibility merely, but responsibility for conduct in the family, the school, and all phases of social relation in which obligation enters as an element. The work is new in plan, and was written to supply a wide-felt want which has not hitherto been met."—*The Popular Science Monthly*.

ROBERTSON.**Mind:** *A Quarterly Review of Psychology and Philosophy.*

Price, \$4.00 per Year.

This new and important periodical, devoted to the science of mind in its modern aspects, has a broad claim upon the thoughtful classes of the community—teachers, professors, physicians, lawyers, clergymen, students, and all interested in the general advance of ideas.

“Mind” is edited by Prof. Croom Robertson, of University College, London, assisted by Prof. Bain and a large number of the ablest psychologists of England; and it will be an organ for the publication of original researches, and a critical record of the progress made in psychology and philosophy in England, on the Continent, and in the United States. Mental phenomena will be taken up on the physiological side, as well as on the side of consciousness, as displayed in the inferior races of mankind, and in the lower animals; in its abnormal phases, as insanity and idiocy; in its relation to language, and in its bearing upon the subject of education. It will also give careful reports of the state and progress of psychological study in the universities of different countries.

MARKOE.**A Treatise on Diseases of the Bones.**

By THOMAS M. MARKOE, M. D.,

Professor of Surgery in the College of Physicians and Surgeons, New York, etc.

WITH NUMEROUS ILLUSTRATIONS.

1 vol., 8vo. Cloth, \$4.50.

This valuable work is a treatise on Diseases of the Bones, embracing their structural changes as affected by disease, their clinical history and treatment, including also an account of the various tumors which grow in or upon them. None of the *injuries* of bone are included in its scope, and no *joint* diseases, excepting where the condition of the bone is a prime factor in the problem of disease. As the work of an eminent surgeon of large and varied experience, it may be regarded as the best on the subject, and a valuable contribution to medical literature.

“The book which I now offer to my professional brethren contains the substance of the lectures which I have delivered during the past twelve years at the college. . . . I have followed the leadings of my own studies and observations, dwelling more on those branches where I had seen and studied most, and perhaps too much neglecting others where my own experience was more barren, and therefore to me less interesting. I have endeavored, however, to make up the deficiencies of my own knowledge by the free use of the materials scattered so richly through our periodical literature, which scattered leaves it is the right and the duty of the systematic writer to collect and to embody in any account he may offer of the state of a science at any given period.”—*Extract from Author's Preface.*

MEYER.

Electricity in its Relations to Practical *Medicine.*

By DR. MORITZ MEYER,
Royal Counsellor of Health, etc.


Translated from the Third German Edition, with Notes and Additions,
A New and Revised Edition,

By WILLIAM A. HAMMOND, M. D.,
Professor of Diseases of the Mind and Nervous System, and of Clinical Medicine, in the Bellevue
Hospital Medical College; Vice-President of the Academy of Mental Sciences, National
Institute of Letters, Arts, and Sciences; late Surgeon-General U. S. A., etc.

1 vol., 8vo. 497 pp. Cloth, \$4.50.

"It is the duty of every physician to study the action of electricity, to become acquainted with its value in therapeutics, and to follow the improvements that are being made in the apparatus for its application in medicine, that he may be able to choose the one best adapted to the treatment of individual cases, and to test a remedy fairly and without prejudice, which already, especially in nervous diseases, has been used with the best results, and which promises to yield an abundant harvest in a still broader domain."—*From Author's Preface.*

SPECIMEN OF ILLUSTRATIONS.



Saxton-Ettinghausen Apparatus.

"Those who do not read German are under great obligations to William A. Hammond, who has given them not only an excellent translation of a most excellent work, but has given us much valuable information and many suggestions from his own personal experience."—*Medical Record.*

"Dr. Moritz Meyer, of Berlin, has been for more than twenty years a laborious and conscientious student of the application of electricity to practical medicine, and the results of his labors are given in this volume. Dr. Hammond, in making a translation of the third German edition, has done a real service to the profession of this country and of Great Britain. Plainly and concisely written, and simply and clearly arranged, it contains just what the physician wants to know on the subject."—*N. Y. Medical Journal.*

"It is destined to fill a want long felt by physicians in this country."—*Journal of Obstetrics*

NIEMEYER.

A Text-Book of Practical Medicine.

With Particular Reference to Physiology and Pathological Anatomy.

By the late Dr. FELIX VON NIEMEYER,

Professor of Pathology and Therapeutics; Director of the Medical Clinic of the University of Tübingen.

Translated from the Eighth German Edition, by special permission of the Author,

By GEORGE H. HUMPHREYS, M. D.,

Late one of the Physicians to the Bureau of Medical and Surgical Relief at Bellevue Hospital for the Out-door Poor; Fellow of the New York Academy of Medicine, etc.,

and

CHARLES E. HACKLEY, M. D.,

One of the Physicians to the New York Hospital; one of the Surgeons to the New York Eye and Ear Infirmary; Fellow of the New York Academy of Medicine, etc.

Revised Edition. 2 vols., 8vo. 1,528 pp. Cloth, \$9.00; Sheep, \$11.00.

The author undertakes, first, to give a picture of disease which shall be as lifelike and faithful to nature as possible, instead of being a mere theoretical scheme; secondly, so to utilize the more recent advances of pathological anatomy, physiology, and physiological chemistry, as to furnish a clearer insight into the various processes of disease.

The work has met with the most flattering reception and deserved success; has been adopted as a text-book in many of the medical colleges both in this country and in Europe; and has received the very highest encomiums from the medical and secular press.

"It is comprehensive and concise, and is characterized by clearness and originality."—*Dublin Quarterly Journal of Medicine.*

"Its author is learned in medical literature; he has arranged his materials with care and judgment, and has thought over them."—*The Lancet.*

"As a full, systematic, and thoroughly practical guide for the student and physician, it is not excelled by any similar treatise in any language."—*Appletons' Journal.*

"The author is an accomplished pathologist and practical physician; he is not only capable of appreciating the new discoveries, which during the last ten years have been unusually numerous and important in scientific and practical medicine, but, by his clinical experience, he can put these new views to a practical test, and give judgment regarding them."—*Edinburgh Medical Journal.*

"From its general excellence, we are disposed to think that it will soon take its place among the recognized text-books."—*American Quarterly Journal of Medical Sciences.*

"The first inquiry in this country regarding a German book generally is, 'Is it a work of practical value?' Without stopping to consider the justness of the American idea of the 'practical,' we can unhesitatingly answer, 'It is!'"—*New York Medical Journal.*

"The author has the power of sifting the tares from the wheat—a matter of the greatest importance in a text-book for students."—*British Medical Journal.*

"Whatever exalted opinion our countrymen may have of the author's talents of observation and his practical good sense, his text-book will not disappoint them, while those who are so unfortunate as to know him only by name, have in store a rich treat."—*New York Medical Record.*

NEUMANN. Hand-Book of Skin Diseases.

By DR. ISIDOR NEUMANN,
Lecturer on Skin Diseases in the Royal University of Vienna.

Translated from advanced sheets of the second edition, furnished by the Author; with Notes,

By LUCIUS D. BULKLEY, A. M., M. D.,

Surgeon to the New York Dispensary, Department of Venereal and Skin Diseases; Assistant to the Skin Clinic of the College of Physicians and Surgeons, New York; Member of the New York Dermatological Society, etc., etc.

1 vol., 8vo. About 450 pages and 66 Woodcuts. Cloth, \$4.00.

Prof. Neumann ranks second only to Hebra, whose assistant he was for many years and his work may be considered as a fair exponent of the German practice of Dermatology. The book is abundantly illustrated with plates of the histology and pathology of the skin. The translator has endeavored, by means of notes from French, English, and American sources, to make the work valuable to the student as well as to the practitioner.

"It is a work which I shall heartily recommend to my class of students at the University of Pennsylvania, and one which I feel sure will do much toward enlightening the profession on this subject."—*Louis A. Duhring.*

"I know it to be a good book, and I am sure that it is well translated; and it is interesting to find it illustrated by references to the views of co-laborers in the same field."—*Erasmus Wilson.*

"So complete as to render it a most useful book of reference."—*T. McCall Anderson.*

"There certainly is no work extant which deals so thoroughly with the Pathological Anatomy of the Skin as does this hand-book."—*N. Y. Medical Record.*

"The original notes by Dr. Bulkley are very practical, and are an important adjunct to the text. . . . I anticipate for it a wide circulation."—*Silas Durkee, Boston.*

"I have already twice expressed my favorable opinion of the book in print, and am glad that it is given to the public at last."—*James C. White, Boston.*

"More than two years ago we noticed Dr. Neumann's admirable work in its original shape; and we are therefore absolved from the necessity of saying more than to repeat our strong recommendation of it to English readers."—*Practitioner.*

NEFTEL. Galvano-Therapeutics. *The Physiological and Therapeutical Action of the Galvanic Current upon the Acoustic, Optic, Sympathetic, and Pneumogastric Nerves.*

By WILLIAM B. NEFTEL.

1 vol., 12mo. 161 pp. Cloth, \$1.50.

This book has been republished at the request of several aural surgeons and other professional gentlemen, and is a valuable treatise on the subjects of which it treats. Its author, formerly visiting physician to the largest hospital of St. Petersburg, has had the very best facilities for investigation.

"This little work shows, as far as it goes, full knowledge of what has been done on the subjects treated of, and the author's practical acquaintance with them."—*New York Medical Journal.*

"Those who use electricity should get this work, and those who do not should peruse it to learn that there is one more therapeutical agent that they could and should possess."—*The Medical Investigator.*

NIGHTINGALE.

Notes on Nursing: *What it is, and what it is not.*

BY FLORENCE NIGHTINGALE.

1 vol., 12mo. 140 pp. Cloth, 75 cents.

Every-day sanitary knowledge, or the knowledge of nursing, or, in other words, of how to put the constitution in such a state as that it will have no disease or that it can recover from disease, takes a higher place. It is recognized as the knowledge which every one ought to have—distinct from medical knowledge, which only a profession can have.

PAGET.

Clinical Lectures and Essays.

BY SIR JAMES PAGET, BART.,

F. R. S., D. C. L., Oxon., LL. D., Cantab.; Sergeant-Surgeon Extraordinary to her Majesty the Queen; Surgeon to H. R. H. the Prince of Wales; Consulting Surgeon to St. Bartholomew's Hospital.

EDITED BY HOWARD MARSH,

Assistant Surgeon to St. Bartholomew's Hospital, and to the Hospital for Sick Children.

1 vol., 8vo. Cloth. Price, \$5.00.

CONTENTS.

THE VARIOUS RISKS OF OPERATIONS—THE CALAMITIES OF SURGERY—STAMMERING WITH OTHER ORGANS THAN THOSE OF SPEECH—CASES THAT BONE-SETTERS CURE—STRANGULATED HERNIA—CHRONIC PYÆMIA—NERVOUS MIMICRY—TREATMENT OF CARBUNCLE—SEXUAL HYPOCHONDRIASIS—GOUTY PHLEBITIS—RESIDUAL ABSCESS—DISSECTION—POISONS—QUIET NECROSIS—SENILE SCROFULA—SCARLET FEVER AFTER OPERATIONS—NOTES FOR THE STUDY OF SOME CONSTITUTIONAL DISEASES—NOTES—INDEX.

PEREIRA.

Dr. Pereira's Elements of *Materia Medica and Therapeutics*. *Abridged and adapted for the Use of Medical and Pharmaceutical Practitioners and Students, and comprising all the Medicines of the British Pharmacopœia, with such others as are frequently ordered in Prescriptions, or required by the Physician.*

EDITED BY ROBERT BENTLEY AND THEOPHILUS REDWOOD.

New Edition. Brought down to 1872. 1 vol., Royal 8vo. Cloth, \$7.00; Sheep, \$8.00.

PEASLEE.

Ovarian Tumors; *Their Pathology, Diagnosis, and Treatment, with Reference especially to Ovariectomy.*

By E. R. PEASLEE, M.D.,

Professor of Diseases of Women in Dartmouth College; one of the Consulting Physicians to the New York State Woman's Hospital; formerly Professor of Obstetrics and Diseases of Women in the New York Medical College; Corresponding Member of the Obstetrical Society of Berlin, etc.

1 vol., 8vo. Illustrated with many Woodcuts, and a Steel Engraving of Dr. E. McDowell, the "Father of Ovariectomy." Price, Cloth, \$5.00.

This valuable work, embracing the results of many years of successful experience in the department of which it treats, will prove most acceptable to the entire profession; while the high standing of the author and his knowledge of the subject combine to make the book the best in the language. It is divided into two parts: the first treating of Ovarian Tumors, their anatomy, pathology, diagnosis, and treatment, except by extirpation; the second of Ovariectomy, its history and statistics, and of the operation. Fully illustrated, and abounding with information, the result of a prolonged study of the subject, the work should be in the hands of every physician in the country.

The following are some of the opinions of the press, at home and abroad, of this great work, which has been justly styled, by an eminent critic, "*the most complete medical monograph on a practical subject ever produced in this country.*"

"His opinions upon what others have advised are clearly set forth, and are as interesting and important as are the propositions he has himself to advance; while, there are a freshness, a vigor, an authority about his writing, which great practical knowledge alone can confer."—*The Lancet*.

"Both Wells's and Peaslee's works will be received with the respect due to the great reputation and skill of their authors. Both exist not only as masters of their art, but as clear and graceful writers. In either work the student and practitioner will find the fruits of rich experience, of earnest thought, and of steady, well-balanced judgment. As England is proud of Wells, so may America well be proud of Peaslee, and the great world of science may be proud of both."—*British Medical Journal*.

"This is an excellent work, and does great credit to the industry, ability, science, and learning of Dr. Peaslee. Few works issue from the medical press so complete, so exhaustively learned, so imbued with a practical tone, without losing other substantial good qualities."—*Edinburgh Medical Journal*.

"In closing our review of this work, we cannot avoid again expressing our appreciation of the thorough study, the careful and honest statements, and candid spirit, which characterize it. For the use of the student we should give the preference to Dr. Peaslee's work, not only from its completeness, but from its more methodical arrangement."—*American Journal of Medical Sciences*.

"Dr. Peaslee brings to the work a thoroughness of study, a familiarity with the whole field of history, physiology, pathology, and practical gynecology, not excelled, perhaps, by those of any man who ever performs the operation."—*Medical Record*.

"We deem its careful perusal indispensable to all who would treat ovarian tumors with a good conscience."—*American Journal of Obstetrics*.

"It shows prodigal industry, and embodies within its five hundred and odd pages pretty much all that seems worth knowing on the subject of ovarian diseases."—*Philadelphia Medical Times*.

"Great thoroughness is shown in Dr. Peaslee's treatment of all the details of this very admirable work."—*Boston Medical and Surgical Journal*.

"Indispensable to the American student of gynecology."—*Pacific Medical and Surgical Journal*.

Reports. *Bellevue and Charity Hospital Reports for 1870, containing valuable contributions from*

ISAAC E. TAYLOR, M.D., AUSTIN FLINT, M.D., LEWIS A. SATRE, M.D., WILLIAM A. HAMMOND, M.D., T. GAILLARD THOMAS, M.D., FRANK H. HAMILTON, M.D., and others.

1 vol., 8vo. Cloth, \$4.00.

"These institutions are the most important, as regards accommodations for patients and variety of cases treated, of any on this continent, and are surpassed by but few in the world. The gentlemen connected with them are acknowledged to be among the first in their profession, and the volume is an important addition to the professional literature of this country."—*Psychological Journal*.

RICHARDSON.

Diseases of Modern Life.

BY DR. B. W. RICHARDSON, F. R. S.

1 vol., 12mo. \$2.00.

PART THE FIRST.—PHENOMENA OF DISEASE, INCIDENTAL AND GENERAL.

- Chap. I.—Natural Life to Natural Death. Euthanasia.
 “ II.—Phenomena of Disease, Classification and Distribution.
 “ III.—Disease Antecedent to Birth.
 “ IV.—External Origins and Causes of Disease.
 “ V.—Phenomena of Disease, from Causes External and Uncontrollable.
 “ VI.—Phenomena of Disease, from Causes External and Communicable.
 “ VII.—Phenomena of Disease, incidental to Old Age and Natural Decay.

PART THE SECOND.—PHENOMENA OF DISEASE, INDUCED AND SPECIAL.

- Chap. I.—Definition and Classification of Induced Diseases.
 “ II.—Disease from Worry and Mental Strain (Broken Heart).
 “ III.—Disease from Worry and Mental Strain, continued (Paralysis).
 “ IV.—Disease from Physical Strain.
 “ V.—Disease from Combined Physical and Mental Strain.
 “ VI.—Disease from the Influence of the Passions.
 “ VII.—Disease from Alcohol. Physiological Proem.
 “ VIII.—Phenomena of Disease from Alcohol. The Functional Type.
 “ IX.—Organic Disease from Alcohol.
 “ X.—Disease from Tobacco. Physiological Phenomena.
 “ XI.—Disease from Tobacco, continued (of the Heart and Lungs).
 “ XII.—Disease from Tobacco, continued (of the Brain and Nervous System).
 “ XIII.—Disease from the Use of Narcotics (from Opium, Chloral, and Absinthe).
 “ XIV.—Disease from Misuse of Foods.
 “ XV.—Disease incident to some Occupations.
 “ XVI.—Disease from Late Hours and Deficient Sleep.
 “ XVII.—Disease from Imperfect Supply of Air.
 “ XVIII.—Disease from Imitation and Moral Contagion.

PART THE THIRD.—SUMMARY OF PRACTICAL APPLICATIONS.

STEINER.

Compendium of Children's Diseases.

A Hand-book for Practitioners and Students.

By Dr. JOHANN STEINER,

Professor of the Diseases of Children in the University of Prague, and Physician to the Francis-Joseph Hospital for Sick Children.

Translated from the Second German Edition by Lawson Tait, F. R. C. S.,

Surgeon to the Birmingham Hospital for Women; Consulting Surgeon to the West Bromwich Hospital; Lecturer on Physiology at the Midland Institute.

1 vol., 8vo. Cloth, \$3.50.

TRANSLATOR'S PREFACE.

Dr. Steiner's book has met with such marked success in Germany that a second edition has already appeared, a circumstance which has delayed the appearance of its English form, in order that I might be able to give his additions and corrections.

In Germany the use of the metric system has not yet entirely superseded the local measures; but it is rapidly doing so, as in England. I have, therefore, rendered all thermometric observations in the Centigrade scale, and all measurements in centi- and millimètres.

I have added as an Appendix the "Rules for Management of Infants" which have been issued by the staff of the Birmingham Sick Children's Hospital, because I think that they have set an example by freely distributing these rules among the poor for which they cannot be sufficiently commended, and which it would be wise for other sick children's hospitals to follow.

I have also added a few notes, chiefly, of course, relating to the surgical ailments of children.

BIRMINGHAM, *October*, 1874.

LAWSON TAIT.

STROUD.

The Physical Cause of the Death of Christ, and its Relations to the Principles and Practice of Christianity.

By WILLIAM STROUD, M. D.

With a Letter on the Subject, by Sir James Y. Simpson, Bart., M. D.

1 vol., 12mo. 422 pp. Cloth, \$2.00.

This important and remarkable book is, in its own place, a masterpiece, and will be considered as a standard work for many years to come.

"The principal point insisted on is, that the death of Christ was caused by rupture or laceration of the heart. Sir James Y. Simpson, who had read the author's treatise and various comments on it, expressed himself very positively in favor of the views maintained by Dr. Stroud."
—*Psychological Journal*.

SIMPSON.

The Posthumous Works of Sir James Young Simpson, Bart., M. D. In Three Volumes.

Volume I.—*Selected Obstetrical and Gynecological Works of Sir James Y. Simpson, Bart., M. D., D. C. L.*, late Professor of Midwifery in the University of Edinburgh. Containing the substance of his Lectures on Midwifery. Edited by J. WATT BLACK, A. M., M. D., Member of the Royal College of Physicians, London; Physician-Accoucheur to Charing Cross Hospital, London; and Lecturer on Midwifery and Diseases of Women and Children in the Hospital School of Medicine.

1 vol., 8vo. 852 pp. Cloth, \$3.00.

This volume contains all the more important contributions of Sir James Y. Simpson to the study of obstetrics and diseases of Women, with the exception of his clinical lectures on the latter subject, which will shortly appear in a separate volume. This first volume contains many of the papers reprinted from his Obstetric Memoirs and Contributions, and also his Lecture Notes, now published for the first time, containing the substance of the practical part of his course of midwifery. It is a volume of great interest to the profession, and a fitting memorial of its renowned and talented author.

"To many of our readers, doubtless, the chief of the papers it contains are familiar. To others, although probably they may be aware that Sir James Simpson has written on the subjects, the papers themselves will be new and fresh. To the first class we would recommend this edition of Sir James Simpson's works, as a valuable volume of reference; to the latter, as a collection of the works of a great master and improver of his art, the study of which cannot fail to make them better prepared to meet and overcome its difficulties."—*Medical Times and Gazette*.

Volume II.—*Anæsthesia, Hospitalism, etc.* Edited by Sir WALTER SIMPSON, Bart.

1 vol., 8vo. 560 pp. Cloth, \$3.00.

"We say of this, as of the first volume, that it should find a place on the table of every practitioner; for, though it is patchwork, each piece may be picked out and studied with pleasure and profit."—*The Lancet (London)*.

Volume III.—*The Diseases of Women.* Edited by ALEX. SIMPSON, M. D., Professor of Midwifery in the University of Edinburgh.

1 vol., 8vo. Cloth, \$3.00.

One of the best works on the subject extant. Of inestimable value to every physician.

SWETT.

A Treatise on the Diseases of the Chest. Being a Course of Lectures delivered at the New York Hospital.

By JOHN A. SWETT, M. D.,

Professor of the Institutes and Practice of Medicine in the New York University; Physician to the New York Hospital; Member of the New York Pathological Society.

1 vol., 8vo. 587 pp. \$3.50.

Embodied in this volume of lectures is the experience of ten years in hospital and private practice.

SAYRE.

A Practical Manual on the Treatment of Club-Foot.

By LEWIS A. SAYRE, M. D.,

Professor of Orthopedic Surgery in Bellevue Hospital Medical College; Surgeon to Bellevue and Charity Hospitals, etc.

1 vol., 12mo. New and Enlarged Edition. Cloth, \$1.00.

"The object of this work is to convey, in as concise a manner as possible, all the practical information and instruction necessary to enable the general practitioner to apply that plan of treatment which has been so successful in my own hands."—*Preface*.

"The book will very well satisfy the wants of the majority of general practitioners, for whose use, as stated, it is intended."—*New York Medical Journal*.

SMITH.

On Foods.

By EDWARD SMITH, M. D., LL. B., F. R. S.,

Fellow of the Royal College of Physicians of London, etc., etc.

1 vol., 12mo. Cloth. Price, \$1.75.

"Since the issue of the author's work on 'Practical Dietary,' he has felt the want of another, which would embrace all the generally-known and less-known foods, and contain the latest scientific knowledge respecting them. The present volume is intended to meet this want, and will be found useful for reference, to both scientific and general readers. The author extends the ordinary view of foods, and includes water and air, since they are important both in their food and sanitary aspects.

SCHROEDER.

A Manual of Midwifery. Including the Pa- thology of Pregnancy and the Puerperal State.

By Dr. KARL SCHROEDER,

Professor of Midwifery and Director of the Lying-in Institution in the University of Erlangen.

TRANSLATED FROM THE THIRD GERMAN EDITION

By CHARLES H. CARTER, B. A., M. D., B. S. London,

Member of the Royal College of Physicians, London, and Physician Accoucheur to St. George's, Hanover Square, Dispensary.

With Twenty-six Engravings on Wood. 1 vol., 8vo. Cloth.

"The translator feels that no apology is needed in offering to the profession a translation of Schroeder's Manual of Midwifery. The work is well known in Germany and extensively used as a text-book; it has already reached a third edition within the short space of two years, and it is hoped that the present translation will meet the want, long felt in this country, of a manual of midwifery embracing the latest scientific researches on the subject.

TILT.

A Hand-Book of Uterine Therapeu- tics and of Diseases of Women.

By EDWARD JOHN TILT, M. D.,

Member of the Royal College of Physicians; Consulting Physician to the Farringdon General Dispensary; Fellow of the Royal Medical and Chirurgical Society, and of several British and foreign societies.

1 vol., 8vo. 345 pp. Cloth, \$3.50.

Second American edition, thoroughly revised and amended.

"In giving the result of his labors to the profession the author has done a great work. Our readers will find its pages very interesting, and, at the end of their task, will feel grateful to the author for many very valuable suggestions as to the treatment of uterine diseases."—*The Lancet*.

"Dr. Tilt's 'Hand-Book of Uterine Therapeutics' supplies a want which has often been felt. . . It may, therefore, be read not only with pleasure and instruction, but will also be found very useful as a book of reference."—*The Medical Mirror*.

"Second to none on the therapeutics of uterine disease."—*Journal of Obstetrics*.

VAN BUREN AND KEYES

A Practical Treatise on the Surgical

Diseases of the Genito-Urinary Organs, including Syphilis. Designed as a Manual for Students and Practitioners. With Engravings and Cases.

BY W. H. VAN BUREN, A. M., M. D.,

Professor of Principles of Surgery, with Diseases of the Genito-Urinary System and Clinical Surgery, in Bellevue Hospital Medical College; Consulting Surgeon to the New York Hospital, the Charity Hospital, etc.; and

E. L. KEYES, A. M., M. D.,

Professor of Dermatology in Bellevue Hospital Medical College; Surgeon to the Charity Hospital, Venereal Diseases; Consulting Dermatologist to the Bureau of Out-Door Relief, Bellevue Hospital, etc.

1 vol., 8vo. Cloth, \$5.00; Sheep, \$6.00.

This work is really a compendium of, and a book of reference to, all modern works treating in any way of the surgical diseases of the genito-urinary organs. At the same time, no other single book contains so large an array of *original* facts concerning the class of diseases with which it deals. These facts are largely drawn from the extensive and varied experience of the authors.

Many important branches of genito-urinary diseases, as the cutaneous maladies of the penis and scrotum, receive a thorough and exhaustive treatment that the professional reader will search for elsewhere in vain.

The work is elegantly and profusely illustrated, and enriched by fifty-five original cases, setting forth obscure and difficult points in diagnosis and treatment.

"The first part is devoted to the Surgical Diseases of the Genito-Urinary Organs; and part second treats of Chancroid and Syphilis. The authors 'appear to have succeeded admirably in giving to the world an exhaustive and reliable treatise on this important class of diseases.'"—*Northwestern Medical and Surgical Journal*.

"It is a most complete digest of what has long been known, and of what has been more recently discovered, in the field of syphilitic and genito-urinary disorders. It is perhaps not an exaggeration to say that no single work upon the same subject has yet appeared, in this or any foreign language, which is superior to it."—*Chicago Medical Examiner*.

"The commanding reputation of Dr. Van Buren in this specialty and of the great school and hospital from which he has drawn his clinical materials, together with the general interest which attaches to the subject-matter itself, will, we trust, lead very many of those for whom it is our office to cater, to possess themselves at once of the volume and form their own opinions of its merit."—*Atlanta Medical and Surgical Journal*.

Lectures upon Diseases of the Rectum.

*Delivered at the Bellevue Hospital Medical College.
Session of 1869-'70.*

BY W. H. VAN BUREN, M. D.,

1 vol., 12mo. 164 pages. Cloth, \$1.50.

"It seems hardly necessary to more than mention the name of the author of this admirable little volume in order to insure the character of his book. No one in this country has enjoyed greater advantages, and had a more extensive field of observation in this specialty, than Dr. Van Buren, and no one has paid the same amount of attention to the subject. . . . Here is the experience of years summed up and given to the professional world in a plain and practical manner."—*Psychological Journal*.

VOGEL.

A Practical Treatise on the Diseases
of Children. Second American from the Fourth
German Edition. Illustrated by Six Lithographic
Plates.

By ALFRED VOGEL, M. D.,

Professor of Clinical Medicine in the University of Dorpat, Russia.

TRANSLATED AND EDITED BY

H. R A P H A E L, M. D.,

Late House Surgeon to Bellevue Hospital; Physician to the Eastern Dispensary for the Diseases
of Children, etc., etc.

1 vol., 8vo. 611 pp. Cloth, \$4.50.

The work is well up to the present state of pathological knowledge; complete without unnecessary prolixity; its symptomatology accurate, evidently the result of careful observation of a competent and experienced clinical practitioner. The diagnosis and differential relations of diseases to each other are accurately described, and the therapeutics judicious and discriminating. All polypharmacy is discarded, and only the remedies which appeared useful to the author commended.

It contains much that must gain for it the merited praise of all impartial judges, and prove it to be an invaluable text-book for the student and practitioner, and a safe and useful guide in the difficult but all-important department of Pædiatrica.

"Rapidly passing to a fourth edition in Germany, and translated into three other languages, America now has the credit of presenting the first English version of a book which must take a prominent, if not the leading, position among works devoted to this class of disease."—*N. Y. Medical Journal*.

"The profession of this country are under many obligations to Dr. Raphael for bringing, as he has done, this truly valuable work to their notice."—*Medical Record*.

"The translator has been more than ordinarily successful, and his labors have resulted in what, in every sense, is a valuable contribution to medical science."—*Psychological Journal*.

"We do not know of a compact text-book on the diseases of children more complete, more comprehensive, more replete with practical remarks and scientific facts, more in keeping with the development of modern medicine, and more worthy of the attention of the profession, than that which has been the subject of our remarks."—*Journal of Obstetrics*.

WALTON.

The Mineral Springs of the United States and Canada, with Analyses and Notes on the Prominent Spas of Europe, and a List of Sea-side Resorts. An enlarged and revised edition.

By GEORGE E. WALTON, M. D.,
Lecturer on Materia Medica in the Miami Medical College, Cincinnati.

Second Edition, revised and enlarged. 1 vol., 12mo. 390 pp., with Maps. \$2.00.

The author has given the analyses of all the springs in this country and those of the principal European spas, reduced to a uniform standard of one wine-pint, so that they may readily be compared. He has arranged the springs of America and Europe in seven distinct classes, and described the diseases to which mineral waters are adapted, with references to the class of waters applicable to the treatment, and the peculiar characteristics of each spring as near as known are given—also, the location, mode of access, and post-office address of every spring are mentioned. In addition, he has described the various kinds of baths and the appropriate use of them in the treatment of disease.

EXTRACTS FROM OPINIONS OF THE PRESS.

“ . . . Precise and comprehensive, presenting not only reliable analyses of the waters, but their therapeutic value, so that physicians can hereafter advise their use as intelligently and beneficially as they can other valuable alterative agents.”—*Sanitarian*.

“ . . . Will tend to enlighten both the profession and the people on this question.”—*N. Y. Medical Journal*.

“ . . . Contains in brief space a vast amount of important and interesting matter, well arranged and well presented. Nearly every physician needs just such a volume ”—*Richmond and Louisville Medical Journal*.

“ . . . Fills this necessity in a scientific and pleasing manner, and can be read with advantage by the physician as well as layman.”—*American Jour. of Obstetrics*.

UNIVERSITY OF VIRGINIA, June 9, 1873.

GENTLEMEN: I have received by mail a copy of Dr. Walton's work on the Mineral Springs of the United States and Canada. Be pleased to accept my thanks for a work which I have been eagerly looking for ever since I had the pleasure of meeting the author in the summer of 1871. He satisfied me that he was well qualified to write a reliable work on this subject, and I doubt not he has met my expectations. Such a work was greatly needed, and, if offered for sale at the principal mineral springs of the country, will, I believe, command a ready sale. Very respectfully yours,

J. L. CABELL, M. D.

WELLS.

Diseases of the Ovaries; *Their Diagnosis and Treatment.*

By T. SPENCER WELLS,

Fellow and Member of Council of the Royal College of Surgeons of England; Honorary Fellow of the King and Queen's College of Physicians in Ireland; Surgeon in Ordinary to the Queen's Household; Surgeon to the Samaritan Hospital for Women; Member of the Imperial Society of Surgery of Paris, of the Medical Society of Paris, and of the Medical Society of Sweden; Honorary Member of the Royal Society of Medical and Natural Science of Brussels, and of the Medical Societies of Pesh and Helsingfors; Honorary Fellow of the Obstetrical Societies of Berlin and Leipzig.

1 vol., 8vo. 478 pp. Illustrated. Cloth, Price, \$4.50.

In 1865 the author issued a volume containing reports of one hundred and fourteen cases of Ovariectomy, which was little more than a simple record of facts. The book was soon out of print, and, though repeatedly asked for a new edition, the author was unable to do more than prepare papers for the Royal Medical and Chirurgical Society, as series after series of a hundred cases accumulated. On the completion of five hundred cases he embodied the results in the present volume, an entirely new work, for the student and practitioner, and trusts it may prove acceptable to them and useful to suffering women.

"Arrangements have been made for the publication of this volume in London on the day of its publication in New York." French and German translations are already in press.

WAGNER.

A Hand - book of Chemical Technology.

By RUDOLPH WAGNER, Ph. D.,

Professor of Chemical Technology at the University of Wurtzburg.

Translated and edited, from the eighth German edition, with extensive additions,

By WILLIAM CROOKES, F. R. S.

With 336 Illustrations. 1 vol., 8vo. 761 pages. Cloth, \$5.00.

Under the head of Metallurgical Chemistry, the latest methods of preparing Iron, Cobalt, Nickel, Copper, Copper Salts, Lead and Tin, and their Salts, Bismuth, Zinc, Zinc Salts, Cadmium, Antimony, Arsenic, Mercury, Platinum, Silver, Gold, Manganates, Aluminum, and Magnesium, are described. The various applications of the Voltaic Current to Electro-Metallurgy follow under this division. The preparation of Potash and Soda Salts, the manufacture of Sulphuric Acid, and the recovery of Sulphur from Soda Waste, of course occupy prominent places in the consideration of chemical manufactures. It is difficult to over-estimate the mercantile value of Mond's process, as well as the many new and important applications of Bisulphide of Carbon. The manufacture of Soap will be found to include much detail. The Technology of Glass, Stone-ware, Limes, and Mortars, will present much of interest to the Builder and Engineer. The Technology of Vegetable Fibres has been considered to include the preparation of Flax, Hemp, Cotton, as well as Paper-making; while the applications of Vegetable Products will be found to include Sugar-boiling, Wine and Beer Brewing, the Distillation of Spirits, the Baking of Bread, the Preparation of Vinegar, the Preservation of Wood, etc.

Dr. Wagner gives much information in reference to the production of Potash from Sugar residues. The use of Baryta Salts is also fully described, as well as the preparation of Sugar from Beet-roots. Tanning, the Preservation of Meat, Milk, etc., the Preparation of Phosphorus and Animal Charcoal, are considered as belonging to the Technology of Animal Products. The Preparation of Materials for Dyeing has necessarily required much space; while the final sections of the book have been devoted to the Technology of Heating and Illumination.

NEW MEDICAL WORKS IN PRESS.

BARTHOLOW'S TREATISE ON THERAPEUTICS.

CONTRIBUTIONS TO REPARATIVE SURGERY, showing its Application to the Treatment of Deformities, produced by Destructive Disease or Injury; Congenital Defects from Arrest or Excess of Development; and Cicatricial Contractions following Burns. Illustrated by Thirty Cases and fine Engravings. By GURDON BUCK, M. D.

ACNE; its Pathology, Etiology, Prognosis, and Treatment. By L. DUNOAN BULKLEY, A. M., M. D., New York Hospital. A monograph of about seventy pages, illustrated, founded on an analysis of two hundred cases of various forms of Acne.

LECTURES ON ORTHOPEDIC SURGERY AND DISEASES OF THE JOINTS, delivered at Bellevue Hospital Medical College during the Winter Session of 1874-1875, by LEWIS A. SAYRE, M. D., Professor of Orthopedic Surgery, Fractures and Dislocations, and Clinical Surgery, in Bellevue Hospital Medical College; Surgeon to Bellevue Hospital; Consulting Surgeon to Charity Hospital; Consulting Surgeon to St. Elizabeth's Hospital; Consulting Surgeon to Northwestern Dispensary; Member of the American Medical Association; Permanent Member of the New York State Medical Society; Fellow of the New York Academy of Medicine; Member of the New York County Medical Society, of the New York Pathological Society, of the Society of Neurology, of the Medico-Legal Society; Honorary Member of the New Brunswick Medical Society; Honorary Member of the Medical Society of Norway; Knight of the Order of Wasa, by His Majesty the King of Sweden, etc., etc. Illustrated by Numerous Wood-Engravings. 1 vol., Svo. Cloth, \$5.00; Sheep, \$6.00.

The publishers take pleasure in announcing that the above work is now passing through the press, and will be ready by the 20th of April, 1876. It is being published at the request of medical gentlemen of the highest standing, in different sections of our country, as well as many abroad, who are anxious to have Dr. Sayre's peculiar views and extended experience in this specialty given to the profession in a plain and practical manner. The book contains the substance of his course of lectures delivered at Bellevue Hospital Medical College, as well as many important cases from his note-book, and from the hospital records. He has also added a number of cases before presented by him to the profession in medical journals, or at the different medical societies, which are considered worthy of permanent record.

The work is enriched by beautiful and excellent illustrations, engraved from original drawings and photographs prepared expressly therefor. The author having enjoyed exceptional opportunities for the study and treatment of these diseases, the results of his labors cannot fail to be of inestimable value to every student and practitioner, and of service to suffering humanity.

Students desiring early copies will please send their orders direct to the publishers,

D. APPLETON & CO., Publishers,

549 & 551 BROADWAY, NEW YORK.

THE NEW YORK MEDICAL JOURNAL.

JAMES B. HUNTER, M. D., Editor.

Published Monthly. Volumes begin in January and July.

"Among the numerous records of Medicine and the collateral sciences published in America, the above Journal occupies a high position, and deservedly so."—*The Lancet (London)*.

"One of the best medical journals, by-the-by, published on the American Continent."—*London Medical Times and Gazette*.

"A very high-class journal."—*London Medical Mirror*.

"The editor and the contributors rank among our most distinguished medical men, and each number contains matter that does honor to American medical literature."—*Boston Journal of Chemistry*.

"Full of valuable original papers, abounding in scientific ability."—*Chicago Medical Times*

"We know no other periodical that we would rather present as a specimen of American skill and intelligence than the NEW YORK MEDICAL JOURNAL."—*Franklin Repository*.

"The NEW YORK MEDICAL JOURNAL, edited by Dr. James B. Hunter, is one of the sterling periodicals of this country. The present editor has greatly improved the work, and evinces a marked aptitude for the responsible duties so well discharged. The contents of this journal are always interesting and instructive; its original matter is often classic in value, and the selected articles are excellent exponents of the progress and truth of medical science."—*Richmond and Louisville Medical Journal*.

Terms, \$4.00 per Annum; or 40 Cents per Number.

THE POPULAR SCIENCE MONTHLY.

Conducted by Prof. E. L. YOUMANS.

Each Number contains 128 pages, with numerous Descriptive and Attractive Illustrations.

Published Monthly. Volumes begin in May and November.

Terms, \$5 per Annum, or Fifty Cents per Number

THE POPULAR SCIENCE MONTHLY was started to promote the diffusion of valuable scientific knowledge, in a readable and attractive form, among all classes of the community, and has thus far met a want supplied by no other periodical in the United States.

The great feature of the magazine is, that its contents are not what science *was* ten or more years since, but what it *is* to-day, fresh from the study, the laboratory, and the experiment: clothed in the language of the authors, inventors, and scientists themselves, which comprise the leading minds of England, France, Germany, and the United States. Among popular articles, covering the whole range of NATURAL SCIENCE, we have the latest thoughts and words of Herbert Spencer, and Professors Huxley, Tyndall, and R. A. Proctor. Since the start, it has proved a gratifying success to every friend of scientific progress and universal education; and those who believed that science could not be made any thing but dry study, are disappointed.

The press all over the land is warmly commending it. We subjoin a few encomiums from those recently given:

"A journal which promises to be of eminent value to the cause of popular education in this country."—*New York Tribune*.

"It is, beyond comparison, the best attempt at journalism of the kind ever made in this country."—*Home Journal*.

"The initial number is admirably constituted."—*Evening Mail*.

"In our opinion, the right idea has been happily hit in the plan of this new monthly."—*Buffalo Courier*.

"Just the publication needed at the present day."—*Montreal Gazette*.

Payment, in all cases, must be made in advance.

Remittances should be made by postal money-order or check to the Publishers,

D. APPLETON & CO.,

549 & 551 Broadway, New York.



COUNTWAY LIBRARY OF MEDICINE

RD

118

B85

RARE BOOKS DEPARTMENT

